



## Effects of Dietary Supplementation of Zinc and $\alpha$ -Tocopheryl Acetate on Performance and Zinc Concentrations in Egg and Tissues of Japanese Quails

Aghaei A<sup>1</sup>, Khosravinia H<sup>1</sup>, Mamuoee M<sup>2</sup>, Azarfar A<sup>1</sup> & Shahriari A<sup>3</sup>

<sup>1</sup>Department of Animal Science, Faculty of Agriculture, Lorestan University, Khorramabad, Iran

<sup>2</sup>Department of Animal Sciences, Faculty of Animal and Food Science, Ramin Agriculture and Natural Resources University, Khuzestan, Iran

<sup>3</sup>Department of Basic Sciences, Faculty of Veterinary Medicine, Chamran University, Ahvaz, Iran

Poultry Science Journal 2017, 5 (1): 57-64

### Keywords

Zinc oxide  
Performance  
Zn enrichment  
Japanese quail  
 $\alpha$ -tocopheryl acetate

### Corresponding author

Heshmatollah Khosravinia  
[khosravi\\_fafa@yahoo.com](mailto:khosravi_fafa@yahoo.com)

### Article history

Received: November 15, 2016  
Revised: January 9, 2017  
Accepted: February 18, 2017

### Abstract

We investigated the effects of dietary supplementation of zinc (ZnO; 0, 40, 80, 120 and 160 mg/kg) and Vit E ( $\alpha$ -tocopheryl acetate; 0 and 40 IU/kg) on egg production, egg quality and Zn content of egg fractions and tissues in Japanese quails. Using a 5 × 2 factorial design, a total of 960 Japanese quails (*Coturnix coturnix japonica*) at day 70 of age were housed in cages and randomly assigned into one of ten experimental treatments, each with four replicates of 24 birds (16 females and eight males per replicate). Egg production was greater ( $P < 0.05$ ) in birds fed diets containing 160 mg/kg of zinc (Zn) than those fed basal diet (control diet), but vitamin E supplementation had no effect on egg production. Quails fed basal diet supplemented with 80 mg/kg Zn showed a significant improvement in their feed conversion ratio compared to the other birds. Birds supplemented with 80, 120 and 180 mg/kg Zn had stronger egg shells than those fed the control diet, while shell thickness was lower in birds supplemented with 0 and 40 mg/kg of Zn ( $P < 0.05$ ). Enrichment of Zn in egg yolk increased when birds received diets supplemented with 80, 120 and 160 mg/kg Zn compare to control group ( $P < 0.05$ ). Supplementation of diet with Zn increased serum concentration of Zn when fed to quails at 120 mg/kg ( $P < 0.05$ ). Thigh muscle, thigh bone, and liver Zn concentrations increased with concentration of Zn supplementation ( $P < 0.05$ ). Vitamin E supplementation had no effects on laying performance, egg shell quality, and Zn concentrations in egg fractions and tissues of Japanese quail.

### Introduction

Quail production has steadily grown over the past few years because of its high profit and low initial investments (Prabakaran, 2003). Quail rearing, like all types of poultry production, could be profitability optimized with scientific feeding and understanding of quail nutrient requirements. Commercial producers have recently shown great interest in examining the effects of supplemental zinc and vitamin E on

quail raising profitability. Such concerns are derived mainly from scarcity of information on quail zinc (Zn) requirements despite numerous reports confirming its promising effects (along with other minerals and vitamins) on growing broilers and laying hens. Zinc is a trace element necessary for normal growth, bone development, feathering and regulation of appetite in all avian species (Batal *et al.*, 2001). Zinc is also involved in many enzymatic and metabolic functions in the

body (Prasad and Kucuk, 2002): activities of several metabolic enzymes drastically decrease in Zn-deficient animals (Prasad and Kucuk 2002). Scaling of skin, especially on feet, and increased mortality have also been observed in severe cases of Zn deficiency in stressed birds (Sahin *et al.*, 2009). In a 24-week trial, supplementing laying hen rations with 48 mg/kg of Zn had no effect on egg production (Supplee *et al.*, 1958). In contrast, Kienholz *et al.* (1961) found that laying hens fed a soy-based diet containing 10 mg/kg of Zn had lower egg production and hatchability than normal until diets were further supplemented with additional Zn. Since most feed ingredients are marginally deficient in Zn, it is of utmost importance to supplement poultry diets with an additional source of Zn. However, Stahl *et al.*, (1986) found that supplementing diets of laying hens with Zn had no effect on egg production, feed intake, and feed conversion ratio (FCR). Japanese quails are particularly sensitive to dietary Zn deficiency, and it has found that Zn is necessary for their normal growth, feathering and normal skeletal development (Koréneková *et al.*, 2005). Therefore, quails may benefit from with Zn supplements.

Poultry cannot synthesize vitamin E. Therefore, their requirements for vitamin E must be fulfilled *via* dietary sources (Chan *et al.*, 1994). Supplementing laying hen diets with vitamin E increased egg production and oxidative stability, and improved the quality of eggs (Cherian *et al.*, 1996a). Sahin *et al.*, (2006) found no effect on body weight, feed intake, and egg weight in Japanese quail supplemented with vitamin E, though egg production increased. Vitamin E, as a biological antioxidant, has been added to animal diets to improve feed efficiency and immune response. Vitamin E supplementation can also improve the quality of meat and eggs, and increase their vitamin E content for the consumers (Sunder *et al.*, 1997). Salgueiro *et al.* (2000) and Kim *et al.* (1998) demonstrated that Zn, which is involved in some biological antioxidant systems, interacts with vitamin E, because vitamin E status was impaired in Zn-deficient animals. Therefore, this study aimed to investigate the effects of dietary supplementation of zinc and vitamin E on production performance and egg quality of breeder Japanese quail.

## Materials and Methods

### Experimental flock and dietary treatments

All procedures used in this study were

approved by Animal Care Committee of Ramin Agriculture and Natural Resources University, Khuzestan, Iran. Nine hundred and sixty 70-d old Japanese quail (*Coturnix coturnix japonica*) were purchased from a commercial breeder flock. The birds were housed in battery cages (60 cm wide × 100 cm long × 28 cm high) equipped with a raised wire floor, and acclimatized to experimental diets and cages for two weeks. The ambient temperature of hen house was kept between 18 to 26°C. The experimental period lasted 70 d, and throughout this period, birds were subjected to a 16L:8D photoschedule. A corn-soybean basal diet was supplemented with Zn as zinc oxide with 74.5% zinc (0, 40, 80, 120 and 160 mg/kg) and vitamin E as DL-  $\alpha$ -tocopheryl acetate (0 and 40 IU/kg), creating ten experimental treatments. The basal diet was formulated to meet or slightly exceed the nutrient requirements of layer Japanese quail recommended by the NRC (1994; Table 1).

The birds were randomly assigned to one of the ten experimental treatments with four replicates of 24 birds each (sixteen females and eight males). Diets (in mash form) and water were offered to the birds *ad libitum* throughout the experimental period.

### Data Collection

Daily egg production and egg weight were recorded for each cage. Feed intake (FI) was measured weekly and was used to calculate feed conversion ratio (FCR; feed intake divided by weight of eggs produced). Egg mass was calculated as egg weight multiplied by percentage of egg production to estimate grams of egg produced per day. On Day 70 of experiment, one female quail from each group was slaughtered using HALAL procedures and liver and thigh muscles were taken for Zn analysis. On the tenth week of the experiment, two eggs were randomly selected from each group and Zn concentration was determined in the egg yolk, egg white and eggshell individually. To determine strength, thickness, and eggshell proportion, six eggs from each treatment were randomly selected on days 42 and 70 of the experiment and were weighed, broken, and proportional weights of yolks and whites were calculated. The residue of whites were wiped from eggshells, and yolk, white, and eggshell were dried for 48 hrs at 60°C.

**Table 1.** Composition and nutrient content (as fed) of the basal diet

Ingredient	% of the diet
Maize	55.5
Soybean meal	24.5
Wheat	4.0
Fishmeal	3.0
Vegetable oil	3.2
Oyster	7.37
Dicalcium phosphate	1.3
Salt	0.35
DL- Methionine	0.23
Lysine	0.05
Vitamin premix <sup>1</sup>	0.25
Mineral premix <sup>2</sup>	0.25
<i>Calculated nutrient composition</i>	
ME (Kcal/kg)	2900
Protein (%)	18
Methionine (%)	0.54
Lysine (%)	1
Calcium (%)	3.1
Available phosphorus (%)	0.45
Methionine + Cystine	0.82
Zinc (mg/kg)	22.45

<sup>1</sup> Vitamin premix per kg contained: vitamin A, 8000 IU; vitamin D3, 2500 IU; vitamin E, 20 IU; Vitamin K, 2 IU, thiamin, 2 mg; riboflavin, 6 mg; pyridoxine, 3 mg; pantothenic acid, 10 mg; folic acid, 1 mg; biotin, 100 µg; niacin, 40 mg and vitamin B12, 10 µg.

<sup>2</sup> Mineral premix supplied the following per kg of diet: manganese, 70 mg; Iron, 40 mg; copper, 10 mg; choline, 200 mg; iodine, 0.4 mg and selenium, 0.3 mg.

To determine Zn content, samples of yolk, white, and eggshell were ashed in a muffle furnace at 450°C for 12 hrs. Ash was then dissolved in 3 M HCl and transferred to a volumetric flask. The Zn concentration in the HCl extract was determined using atomic spectrometry (Analytic Jena, Contra A [300], Germany) (Skřivan *et al.*, 2005). Zinc concentrations in liver and thigh muscle samples were also determined according to procedures described by Skřivan *et al.* (2005).

Left tibia samples were boiled in deionized water for 20 min, cleaned from all soft tissues, and dried at 60°C for 48 hrs. Fat was extracted with 96% ethanol in a glass container for 48 hrs. During the extraction period, ethanol was replaced several times until its color became clear. Thereafter, samples were dried for 48 hrs at 60°C before ashing at 600°C overnight in a muffle furnace. Zn concentrations from these samples were then determined as described by Skřivan *et al.* (2005).

### Statistical analysis

Statistical analyses were carried out using the MIXED procedure of SAS 9.2 (2003). The model used was:

$$Y_{ijkl} = \mu + B_i + Z_j + E_k + (ZE)_{jk} + e_{ijkl}$$

Where  $Y_{ijkl}$  is the dependent variable under examination;  $\mu$  is the population mean for the variable;  $B_i$  is the random effect of block;  $Z_j$  is the fixed Zn ( $j = 5; 0, 40, 80, 120$  and  $160$  mg/kg of diet);  $E_k$  is the fixed effect of vitamin E ( $k = 2; 0$  and  $40$  IU/kg of diet) and  $e_{ijkl}$  is the random error associated with the observation  $ijk$ .

The Fisher's protected least significant difference (LSD) test was used for multiple treatment comparisons using LSMEANS of SAS 9.2 (2003) with letter grouping obtained using SAS pdmix800 macro (Saxton, 1998). Residual analysis was carried out to test the model assumptions using the UNIVARIATE procedure of SAS 9.2 (2003) with NORMAL and PLOT options. For all statistical analyses, significance was declared at  $P < 0.05$ .

## Results and Discussion

### Performance and egg quality

Effects of dietary supplementation of Zn and vitamin E on performance parameters of Japanese quails are shown in Table 2. Egg production was greater in birds fed diets containing 160 mg/kg Zn compared to those fed 0 or 40 mg/kg Zn in diet ( $P < 0.05$ ). Feed intake and egg weight were not affected by dietary treatments. Compared to control birds, quails

supplemented with 80, 120 or 160 mg/kg Zn showed significant improvement in FCR. Supplee *et al.* (1958) found that supplementing diets with 48 mg/kg Zn had no effect on egg production in layers, but Bahakaim *et al.* (2014) observed an improved FCR when layers were fed diets supplemented with 50 or 100 mg/kg of Zn. Sahin and Kucuk (2003) also reported a linear increase in FI and egg production, and improved feed efficiency and egg quality in heat-stressed quails supplemented with 30 or 60 mg/kg of Zn. However, supplementing layer hens with 100 mg/kg Zn-methionine chelate decreased egg production (Lim *et al.*, 2003). Hermayer *et al.* (1977) found that high levels of Zn in diet reduced feed intake and egg production of laying

hens. Broilers fed diets supplemented with different levels of Zn (40, 80 and 120 mg/kg diet) also reduced feed intake as levels of Zn increased (Refaie, 2009). In line with our findings, Shyam Sunder *et al.* (2008) also observed that supplementing broiler diet with Zn had no effect on feed intake. Kucuk *et al.* (2008) reported that FCR and egg production improved when both Zn (30 mg/kg) and pyridoxine (8 mg/kg) were included in a laying hen diet. Also, Kaya *et al.* (2001) observed an improved FCR when laying hens were fed different levels of supplemental Zn (0, 20, 50, 100 and 200 mg/kg diet), and that the best FCR was observed in birds received diets supplemented with 100 mg/kg of Zn.

**Table 2.** Laying performance and egg quality of Japanese quail supplemented with zinc (Zn) and vitamin E

	Zn (mg/kg of diet)	Vitamin E (IU/kg of diet)	Egg productio n (%)	Feed intake (g/bird/da y)	Egg weig ht (g)	FCR <sup>1</sup>	Egg mass (g/d)	Shell strength (kg/cm <sup>2</sup> )	Shell thickne ss (mm)	Egg shell proportio n (%)
Main effects										
	0		89.08 <sup>c</sup>	32.92	13.37	2.46 <sup>a</sup>	11.91	0.42 <sup>b</sup>	0.196 <sup>b</sup>	12.70
	40		89.39 <sup>bc</sup>	32.71	13.43	2.43 <sup>ab</sup>	12.01	0.57 <sup>ab</sup>	0.195 <sup>b</sup>	12.84
	80		92.55 <sup>ab</sup>	31.47	13.39	2.35 <sup>c</sup>	12.39	0.85 <sup>a</sup>	0.217 <sup>a</sup>	13.38
	120		90.25 <sup>abc</sup>	31.85	13.37	2.38 <sup>bc</sup>	12.08	0.70 <sup>a</sup>	0.210 <sup>a</sup>	13.21
	160		93.17 <sup>a</sup>	32.03	13.51	2.37 <sup>bc</sup>	12.59	0.73 <sup>a</sup>	0.209 <sup>a</sup>	13.52
SEM			1.08	0.41	0.11	0.03	0.20	0.09	0.003	0.396
		0	90.87	32.31	13.41	2.41	12.19	0.70	0.203	13.13
		40	90.90	32.09	13.42	2.39	12.20	0.61	0.208	13.52
SEM			0.68	0.26	0.71	0.02	0.12	0.06	0.002	0.396
Zn × Vit E										
	0	0	89.14	32.98	13.35	2.47	11.90	0.45	0.198	12.75
	0	40	89.02	32.86	13.38	2.46	11.91	0.38	0.195	12.65
	40	0	90.28	33.35	13.41	2.49	12.10	0.69	0.194	12.83
	40	40	88.50	32.07	13.46	2.38	11.92	0.45	0.195	12.84
	80	0	92.04	31.51	13.42	2.35	12.35	0.95	0.212	13.51
	80	40	93.07	31.42	13.36	2.35	12.43	0.75	0.222	13.24
	120	0	90.03	31.69	13.20	2.40	11.88	0.79	0.206	13.18
	120	40	90.46	32.01	13.55	2.36	12.27	0.62	0.215	13.23
	160	0	92.87	31.99	13.67	2.34	12.70	0.64	0.205	13.38
	160	40	93.47	32.08	13.36	2.40	12.49	0.83	0.214	13.65
SEM			1.53	0.58	0.16	0.04	0.28	0.13	0.005	0.559
P-value										
	Zn		0.03	0.10	0.87	0.02	0.10	0.02	0.002	13.13
	Vitamin E		0.97	0.56	0.89	0.41	0.93	0.24	0.804	13.12
	Zn × Vit E		0.90	0.69	0.36	0.27	0.82	0.48	0.507	0.250

<sup>1</sup> FCR, feed conversion ratio

<sup>a,b,c</sup> Lsmeans within a column with different superscripts differ ( $P < 0.05$ ).

In the present study, the improved egg production and FCR in birds that received supplemental Zn could be partly attributed to its role in protection of pancreatic tissue against oxidative damage, which may facilitate proper functioning of the pancreas (e.g. secretion of

digestive enzymes), thereby improving digestibility of nutrients. Furthermore, because Zn is involved in metabolism of carbohydrates, proteins, and lipids (MacDonald, 2000; Ibs and Rink, 2003), energy and protein utilization may have been improved in Zn supplemented birds.

However, in contrast to our findings, El-Latif (1999) reported that egg production in layers was not influenced by supplemental Zn. We found no influence of vitamin E supplementation on laying performance and eggshell quality of breeder quails. In contrast, Cherian *et al.* (1996b) showed that adding vitamin E to poultry diets improved oxidative stability, egg production, and egg quality. In this study, the lack of difference in laying performance in birds supplemented with vitamin E compared to control birds may be that the birds reared under the neutral temperature (21-26°C), and that they received adequate vitamin E (20 IU/kg diet) *via* the basal diet.

Birds supplemented with 80, 120 and 180 mg/kg Zn had stronger egg shells than those fed the control diet, while shell thickness was lower in the birds fed basal diet and basal diet supplemented with 40 mg/kg Zn compared with the other treatments ( $P < 0.05$ ). The results of this study agree with the findings of Mabe *et al.* (2003) who reported that dietary supplementation of laying hens with 60 mg/kg Zn had no effect on percentage of eggshell and eggshell index, but improved egg shell strength. Zinc is directly involved in eggshell synthesis as a co-factor of carbonic anhydrase, which provides carbonate ions in magnum during the deposition of albumen and in isthmus for formation of eggshell membranes. Zn is also involved in eggshell formation pathways in uterine cells (Bahakaim *et al.*, 2014).

### Zn concentration in tissues

Effects of dietary supplementation of Zn and vitamin E on Zn concentration in serum, muscle, bone, liver and egg fractions are shown in Table 3. Serum concentrations of Zn increased when quails were fed 120 mg/kg Zn. Thigh muscle, thigh bone, and liver Zn concentrations improved only when birds were supplemented with 80 mg/kg Zn or greater. Supplementation of diet with Zn in concentrations beyond 40 mg/kg increased yolk concentration of Zn ( $P < 0.05$ ) which agreed with the results of Plaimast *et al.* (2008) who found that Zn deposition in egg increased linearly as dietary Zn levels increased. Henry *et al.* (1987) showed tissue concentration of Zn, especially bone, increased with dietary Zn content. Furthermore, as the levels of dietary zinc increased in turkey, zinc content of tibia, beaks, kidneys, liver, testes and feathers also

increased (Vohra *et al.*, 1968). Tissue uptake of Zn in chicken is related to dietary Zn intake (Sandoval *et al.*, 1997). In agreement with our results, it has been observed that feeding chicken with dietary zinc at levels greater than growth requirements increased Zn concentration in plasma and tibia (Pimentel *et al.*, 1991).

In the present study, Zn supplementation had no effect on eggshell and egg white concentrations of Zn. Yang *et al.*, (2004) indicated that Zn content of egg increased by 55.7% and 70.2% when laying hens were fed diets containing 240 and 840 mg/kg Zn, respectively. Mabe *et al.* (2003) also found a significant increase in Zn content of egg yolk when birds were supplemented with 60 mg/kg Zn. Stahl *et al.*, (1988) reported that hens fed diets containing high levels of zinc (1762 or 1861 mg/kg diet) produced eggs containing 57-95% more zinc than those fed a diet containing 26 mg/kg Zn. In the current study, dietary inclusion of 80 mg Zn/kg of diet increased Zn content of egg yolk by approximately 20% compared with the basal diet, indicating a positive correlation between dietary Zn content and that of the yolk content. Unlike egg yolk, dietary Zn did not change Zn contents of eggshell and egg white most likely due to naturally low content of Zn in these fractions. In contrast to our results, Skřivan *et al.* (2005) found no significant differences in Zn content of egg yolk when 80 mg/kg Zn was added to a basal diet fed of laying hens.

When Zn and vitamin E were supplemented together, there were no significant differences in production performance nor Zn concentration in egg, thigh muscle, bone, liver and serum (Tables 2 and 3). Kim *et al.*, (1998) reported that intestinal absorption of vitamin E was influenced by the Zn status in rats. Zn is involved in certain anti-oxidative enzymatic systems which may interact with vitamin E in stressed birds. Vitamin E status can be impaired in Zn-deficient animals (Salgueiro *et al.*, 2000; Kim *et al.*, 1998). Sahin *et al.* (2006) observed a significant interaction between vitamin E and Zn in improved final body weight and feed intake of growing heat-stressed quails. In this experiment, the lack of interaction between Zn and vitamin E supplementation may be attributed to the fact that birds were raised under neutral temperatures (21-26 °C) and therefore, may not have been exposed to oxidative stress.

**Table 3.** Concentrations of Zn in serum (mg/dL), egg, liver and thigh muscle (mg/kg dry matter) of Japanese quail supplement with zinc and vitamin E

	Zn (mg/kg of diet)	Vitamin E (IU/kg of diet)	Zn concentration					liver	Serum
			egg shell	egg yolk	egg white	Thigh muscle	Thigh bone		
Main effects									
	0		4.30	41.65 <sup>c</sup>	8.23	57.51 <sup>b</sup>	115.75 <sup>c</sup>	66.10 <sup>c</sup>	338.5 <sup>b</sup>
	40		4.24	45.31 <sup>bc</sup>	8.96	59.16 <sup>b</sup>	133.92 <sup>bc</sup>	69.28 <sup>bc</sup>	359.75 <sup>b</sup>
	80		4.80	50.05 <sup>ab</sup>	9.86	64.1 <sup>ab</sup>	153.23 <sup>ab</sup>	73.98 <sup>abc</sup>	397.25 <sup>ab</sup>
	120		4.65	54.17 <sup>a</sup>	8.77	63.42 <sup>ab</sup>	161.52 <sup>a</sup>	80.37 <sup>ab</sup>	406.13 <sup>ab</sup>
	160		4.97	54.11 <sup>a</sup>	9.17	69.69 <sup>a</sup>	173.47 <sup>a</sup>	85.46 <sup>a</sup>	467.63 <sup>a</sup>
SEM			0.25	2.59	0.58	2.51	8.19	3.90	24.85
		0	4.63	48.74	8.88	60.99	149.04	73.55	397.80
		40	4.54	49.38	9.11	64.59	146.12	76.52	389.90
SEM			0.16	1.64	0.37	1.59	5.18	2.47	15.72
Zn × Vit E									
	0	0	3.94	41.91	8.23	57.47	121.44	66.74	339.00
	0	40	4.65	41.39	8.24	57.55	121.44	65.46	337.50
	40	0	4.62	45.82	8.96	55.27	125.61	71.65	340.75
	40	40	3.85	44.81	8.96	63.05	142.24	66.92	378.75
	80	0	4.97	47.41	9.32	64.46	149.90	65.81	401.50
	80	40	4.63	52.69	10.40	63.85	156.55	82.14	393.00
	120	0	4.82	51.84	8.98	58.57	170.25	79.76	429.75
	120	40	4.48	56.50	8.57	68.26	152.79	80.98	382.50
	160	0	4.82	56.71	8.96	69.15	177.98	83.79	477.50
	160	40	5.11	51.52	9.39	70.23	168.97	87.12	457.75
SEM			0.35	3.66	0.82	3.55	11.58	5.51	35.15
P-value									
Zn			0.198	0.006	0.398	0.018	0.001	0.010	0.012
Vitamin E			0.691	0.783	0.668	0.120	0.694	0.401	0.725
Zn × Vit E			0.272	0.594	0.913	0.477	0.573	0.393	0.816

<sup>a,b,c</sup> Lsmeans within a column with different superscripts differ ( $P < 0.05$ ).

### Conclusion

Dietary supplementation with Zn beyond 40 mg/kg of diet improved egg production and eggshell strength and thickness in laying Japanese quail. Zn supplementation also enriched Zn content of egg yolk. Zn contents of

thigh muscle and thigh bone, liver, and serum increased only when supplementation concentration was beyond 80 mg/kg of diet. Supplementary vitamin E had no effect on laying performance of Japanese quail.

### References

- Bahakaim ASA, Abdel Magied HA, Osman SMH, Omar AS, AbdelMalak NY & Ramadan NA. 2014. Effect of using different levels and sources of zinc in layer's diets on egg zinc enrichment. *Egyptian Poultry Science*, 34: 39-56.
- Batal AB, Parr TM & Baker DH. 2001. Zinc bioavailability in tetrabasic zinc chloride and the dietary zinc requirement of young chicks fed a soy concentrate diet. *Poultry Science*, 80: 87-90. DOI: 10.1093/ps/80.1.87.
- Chan KM, Decker EA & Feustman C. 1994. Endogenous skeletal muscle antioxidants. *Critical Reviews in Food Science and Nutrition*, 34: 403-426. DOI: 10.1080/10408399409527669
- Cherian G, Wolfe FH & Sim JS. 1996a. Feeding dietary oils with tocopherols: effects on internal qualities of eggs during storage. *Journal of Food Science*, 61: 15-18. DOI: 10.1111/j.1365-2621.1996.tb14716.x
- Cherian G, Wolfe FW & Sim JS. 1996b. Dietary oils with added tocopherols: effects on egg or tissue tocopherols, fatty acids and oxidative stability. *Poultry Science*, 75: 423-431. DOI: 10.3382/ps.0750423
- El-Latif SAA. 1999. Nutritional interrelationships of vitamin E and selenium on laying Japanese

- quail. *Egyptian Journal of Nutrition and Feeds*, 2: 711-718.
- Henry PR, Ammerman CB & Miles RD. 1987. Effect of dietary zinc on tissue mineral concentration as a measure of zinc bioavailability in chicks. *Nutrition Reports International*, 35: 15-23.
- Hermayer KL, Stake PE & Shippe RL. 1977. Evaluation of dietary zinc, cadmium, tin, lead, bismuth and arsenic toxicity in hens. *Poultry Science*, 56: 1721-1722.
- Ibs KH & Rink L. 2003. Zinc - altered immune function. *The Journal of Nutrition*, 133: 1452-1456.
- Kaya S, Umucalilar HD, Haliloglu S & Ipek H. 2001. Effect of dietary vitamin A and zinc on egg yield and some blood parameters of laying hens. *Turkish Journal of Veterinary and Animal Science*, 25: 763-769.
- Kienholz EW, Turk DE, Sunde ML & Hoekstra WG. 1961. Effects of zinc deficiency in the diets of hens. *Journal of Nutrition*, 75: 211-221.
- Kim ES, Noh SK & Koo SI. 1998. Marginal zinc deficiency lowers the lymphatic absorption of  $\alpha$ -tocopherol in rats. *The Journal of Nutrition*, 128: 265-270.
- Koréneková B, Skalická M, Nad' P, Venglovský J & Sály J. 2005. Supplementation of zinc and cadmium on egg quality of Japanese quails. XIIth International Congress, ISAH, Warsaw, Poland. 2: 118-121.
- Kucuk O, Kahraman A, Kurt I, Yildiz N & Onmaz AC. 2008. A combination of zinc and pyridoxine supplementation to the diet of laying hens improves performance and egg quality. *Biological Trace Element Research*, 126: 165-175. DOI: 10.1007/s12011-008-8190-z
- Lim HS & Paik IK. 2003. Effects of supplementary mineral methionine chelates (Zn, Ca, Mn) on the performance and egg shell quality of laying hens. *Asian-Australasian Journal of Animal Sciences*, 16: 1804-1805. DOI: 10.5713/ajas.2003.1804
- Mabe I, Rapp C, Bain MM & Nys Y. 2003. Supplementation of corn soybean meal diet with manganese, copper and zinc from organic or inorganic sources improves egg shell quality in aged laying hens. *Poultry Science*, 82: 1903-1913. DOI: 10.1093/ps/82.12.1903
- MacDonald RS. 2000. The role of zinc in growth and cell proliferation. *The Journal of Nutrition*, 130: 1500-1508.
- National Research Council. 1994. *Nutrient Requirements of Poultry*, 9<sup>th</sup> edition National Academy Press. Washington. D.C. 196 Pages.
- Pimentel JL, Cook ME & Greger JL. 1991. Research note: bioavailability of zinc-methionine for chicks. *Poultry Science*, 70: 1637-1639. DOI: 10.3382/ps.0701637
- Plaimast H, Sirchakwal P, Puwastien P & Kijparkorn S. 2008. Effect of supplementary zinc from organic sources on zinc deposition in eggs and laying performance. *Journal of Veterinary Medicine*, 38: 47-53.
- Prabakaran R. 2003. Good practices in planning and management of integrated commercial poultry production in South Asia. FAO, Pp. 71-77.
- Prasad AS & Kucuk O. 2002. Zinc in cancer prevention. *Cancer and Metastasis Reviews*, 21: 291-295. DOI: 10.1023/A:1021215111729
- Refae MA. 2009. Performance and immunocompetence of broilers as affected by zinc, protein and Phytase supplementation during summer season. Ph.D. Thesis. Department of Animal nutrition. Faculty of Agriculture, Cairo University.
- Sahin K & Kucuk O. 2003. Zinc supplementation alleviates heat stress in Japanese quail. *The Journal of Nutrition*, 9: 2808-2811.
- Sahin K, Onderci M, Sahin N, Gulcu F, Yildiz N, Avci M & Kucuk O. 2006. Responses of quail to dietary Vitamin E and zinc picolinate at different environmental temperatures. *Animal Feed Science and Technology*, 129: 39-48. DOI: 10.1016/j.anifeedsci.2005.11.009
- Sahin K, Sahin N, Kucuk O, Hayirli A & Prasad AS. 2009. Role of dietary zinc in heat stressed poultry: A review. *Poultry Science*, 88: 2176-2183. DOI: 10.3382/ps.2008-00560
- Sahin N, Sahin K, Onderci M, Karatepe M, Smith MO & Kucuk O. 2006. Effects of dietary lycopene and vitamin E on egg production, antioxidant status and cholesterol levels in Japanese quail. *Asian Australasian Journal of Animal Sciences*, 19: 224-230. DOI: 10.5713/ajas.2006.224
- Salgueiro MJ, Zubillaga M, Lysionek A, Sarabia MI, Caro R, De Paoli T, Hager A, Weill R & Boccio J. 2000. Zinc as an essential micronutrient: A review. *Nutrition Research*, 20: 737-755. DOI: 10.1016/S0271-5317(00)00163-9
- Sandoval M, Henry PR, Ammerman CB, Miles

- RD & Littell RC. 1997. Relative bioavailability of supplemental inorganic zinc sources for chicks. *Journal of Animal Science*, 75: 3195-3205. DOI: 10.2527/1997.75123195x
- SAS Institute. 2003. SAS statistics user's guide. Version 9.2. SAS Institute Inc, Cary, NC. USA.
- Saxton AM. 1998. A macro for converting mean separation output to letter groupings in Proc Mixed. Proc. 23rd SAS Users Group Intl., SAS Institute, Cary, NC.
- Shyam Sunder G, Panda AK, Gopinath NCS, Rama Rao SV, Raju MVLN, Reddy M R & Vijay Kumar Ch. 2008. Effects of higher levels of zinc supplementation on performance, mineral availability, and immune competence in broiler chickens. *The Journal of Applied Poultry Research*, 17: 79-86. DOI: 10.3382/japr.2007-00029
- Skřivan M, Skřivanová V & Marounek M. 2005. Effect of dietary zinc, iron and copper in layer feed on distribution of these elements in eggs, liver, excreta, soil, and herbage. *Poultry Science*, 84: 1570-1575. DOI: 10.1093/ps/84.10.1570
- Stahl JL, Cook ME & Greger JL. 1988. Zinc, iron and copper contents of eggs from hens fed varying levels of zinc. *Journal of Food Composition and Analysis*, 1: 309-315. DOI: 10.1016/0889-1575(88)90030-0
- Stahl LJ, Cook ME & Sunde ML. 1986. Zinc supplementation: Its effect on egg production, feed conversion, fertility, and hatchability. *Poultry Science*, 65: 2104-2109. DOI: 10.3382/ps.0652104
- Sunder A, Richter G & Flachowsky G. 1997. Influence of different concentrations of vitamin E in the feed of laying hens on the vitamin E-transfer into the egg. *Proceedings of the Society of Nutrition Physiology*, 6: 114-152.
- Supplee WC, Blamberg DL, Keene OD, Combs GF & Romoser GL. 1958. Observations on zinc supplementation of poultry rations. *Poultry Science*, 37: 1245 (Abstract).
- Vohra P, Gottfredson GD & Kratzer FH. 1968. The Effects of high levels of dietary EDTA, zinc or copper on the mineral contents of some tissue of turkey poults. *Poultry Science*, 4: 1334-1343. DOI: 10.3382/ps.0471334
- Yang LIE, Coa S, Cheng M, Chen L & Chen K. 2004. Effects of iron, zinc, iodine and selenium levels in rations on activities of metabolic enzymes of layers and egg quality. *Journal of Huazhong Agricultural University*, 51: 352-413.