
Flavonoids as Dietary Additives in Laying Hens: A Meta-analysis of Production Performance, Egg Quality, Liver, and Antioxidant Enzyme Profile

Tri Rachmanto Prihambodo^{1,2}  Muhammad Miftakhus Sholikin³  Nahrowi Nahrowi⁴ 
Irmanida Batubara^{5,6}  Desianto Budi Utomo⁷  & Anuraga Jayanegara⁴ 

¹ Graduate School of Nutrition and Feed Science, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia

² Faculty of Animal Science, Jenderal Soedirman University, Purwokerto 53122, Indonesia

³ National Research and Innovation Agency of Indonesia, Jakarta 10340, Indonesia

⁴ Department of Nutrition and Feed Technology, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia

⁵ Department of Chemistry, Faculty of Mathematics and Natural Sciences, IPB University, Bogor 16680, Indonesia

⁶ Tropical Biopharmaca Research Center (Trop BRC), IPB University, Bogor 16128, Indonesia

⁷ Charoen Pokphand Group Co., Ltd., Jakarta 14430, Indonesia

Poultry Science Journal 2022, 10(1): 27-34

Abstract

In this study, a meta-analysis was employed to evaluate the effects of dietary flavonoids on laying hens. A database of flavonoids was developed from published journal papers in which flavonoids were supplemented to laying hens at various doses and then monitored for production performance, egg quality, liver condition, and antioxidant enzyme profile. A total of 22 journal papers were included in the database, and statistically analyzed using a mixed-model methodology in which random effects consisted of the different studies and fixed effects were the dosages of flavonoids. The model statistics were p-value and Akaike information criterion. The significance of an effect was assumed at $P < 0.05$. The results showed that dietary addition of flavonoids linearly decreased ($P < 0.001$) feed intake. The laying rate was not affected by the addition of flavonoids, while the FCR tended to linearly decrease ($P < 0.1$). The addition of flavonoids linearly reduced egg and yolk cholesterol (both at $P < 0.001$), and linearly increased egg mass ($P < 0.001$), egg strength ($P < 0.05$) and yolk color ($P < 0.001$). Further, a linear decreasing response was observed in the liver triglyceride, and cholesterol concentrations (both at $P < 0.001$) after flavonoid supplementation. Superoxide dismutase showed a quadratic response ($P < 0.01$) following the addition of flavonoids. In conclusion, dietary supplementation of flavonoids is able to favourably modulate production performance and egg quality of laying hens.

Keywords

Laying hen
Flavonoid
Feed intake
Meta-analysis
Superoxide dismutase

Corresponding author

Anuraga Jayanegara
anuragaja@apps.ipb.ac.id

Article history

Received: August 7, 2021
Revised: April 21, 2022
Accepted: May 21, 2022

Introduction

Secondary metabolites, commonly referred to as phytochemicals, are the naturally present complex compounds in plants. These compounds have various benefits for the plants, animals humans, and the environment. Some phytochemicals, however, have anti-nutritional properties that hinder digestion and utilization of nutrients in the digestive tract of animals (Takahama and Hirota, 2018; Panda and Shinde, 2017). The phytochemicals classified as anti-nutrients include flavonoids, tannins, saponins and alkaloids (Kunatsa *et al.*, 2020). In recent years, these

anti-nutritional groups have been avoided due to reduced nutrient absorption in livestock. However, these compounds have become an exciting topic for research because phytochemicals can be used as antibacterial (Prihambodo *et al.*, 2019), anti-tumour, anti-inflammatory, and antiviral substances in an appropriate dosage (Raina *et al.*, 2014)

Flavonoids are one of the largest groups of phytochemicals in plants with numerous benefits. They are often referred to as health-promoting biomolecules and functional ingredients because of their role in improving health and prevention of

chronic diseases. They have often been used in recent decades in livestock and human nutrition (Kamboh *et al.*, 2015). A previous meta-analysis (Prihambodo *et al.*, 2020) has reported positive effects of flavonoids on the performance parameters of broiler chickens. Other authors also reported beneficial effects of flavonoids administration in poultry, such as increasing performance in quails (Juráni *et al.*, 2008) and lowering of cholesterol in ducks (Tugiyanti & Susanti, 2018).

Regarding the effects of flavonoids on laying hens, few studies (increasing growth performance, quality of eggs and enhancing gut morphology and functionality) have been conducted, but there is no study attempting to summarize their results. The meta-analysis is used for statistically combining the results of studies from various literatures in which all influencing factors were calculated (St-Pierre, 2001; Sauviant *et al.* 2008). Therefore, the present study aimed to evaluate the effects of dietary flavonoids administration on production performance, egg quality, liver, and antioxidant enzyme profile in laying hens using a meta-analysis approach.

Materials and Methods

Database development

A database was constructed from published journal papers reporting flavonoids as feed additives in

laying hens. The documents were collected using various searchplatforms such as Scopus, Science Direct, and Google Scholar. The parameters integrated in the database were including production parameters (final body weight, laying rate, feed conversion rate [FCR], and feed intake), egg quality (egg weight, shape index, Haugh unit, egg mass, egg thickness, egg strength, egg cholesterol, shell weight, shell percentage, shell strength, albumen mass, albumen index, albumen height, yolk color, yolk mass, egg mass and yolk cholesterol), liver profile (weight, triglyceride and cholesterol concentrations) and antioxidant enzyme profile (superoxide dismutase, catalase and glutathione peroxidase). The following requirements were used for the inclusion of articles into the database: (a) in vivo studies carried on laying hens; (b) flavonoids were as feed additives, and (c) the articles were published in the English language. Based on the title and abstract screenings, 34 articles met the inclusion criteria. After further full-text evaluation, the final database consisted of 21 studies with 153 treatments, as presented in Table 1. The laying hens included in the database were Hessian, Lohmann Brown, Leghorn, Lohmann White, Hy-line White, ISA Brown, White Leghorn, and Hy-line Brown. In tabulating data into the database, data relating to the parameters were converted into the same measurement units to facilitate further analyses.

Table 1. Studies included in the meta-analysis

No	Reference	Layer Type	Period (weeks)	Flavonoid Type	Dosage (g kg ⁻¹)
1	Galal <i>et al.</i> (2008)	Hy-line White	46-50	Quercetin	0.0.15
2	Lien <i>et al.</i> (2008)	Leghorn	30-42	Hesperitin	0-5
3	Deng <i>et al.</i> (2010)	Hy-line Brown	28-36	Quercetin	0-0.034
4	Ariana <i>et al.</i> (2011)	Hy-line White	45-53	Naringenin	0-5
5	Ting <i>et al.</i> (2011)	White Leghorn	26-36	Hesperitin & naringenin	0-4
6	Liu <i>et al.</i> (2013)	Hessian	39-47	Quercetin	0-0.6
7	Goliomytis <i>et al.</i> (2014)	Lohmann White	26-30	Hesperidin	0-3
8	Liu <i>et al.</i> (2014)	Hessian	28-36	Quercetin	0-0.0619
9	Lokhande <i>et al.</i> (2014)	Hy-line Brown	45-53	Quercetin & hesperdin	0-7.5
10	Simitzis <i>et al.</i> (2014)	Lohmann Brown	28-36	Hesperidin & naringenin	0-1.5
21	Ying <i>et al.</i> (2015)	Hessian	39-47	Quercetin	0-0.6
11	Damaziak <i>et al.</i> (2017)	ISA Brown	16-32	Quercetin	0-0.032
12	El-Tarabany (2018)	Tetra Brown	58-63	Quercetin	0-0.2
13	Iskender <i>et al.</i> (2017)	Lohmann White	28-36	Hesperidin, naringenin & quercetin	0-0.5
14	Li <i>et al.</i> (2017)	Lohmann Brown	72-79	Hesperidin, naringenin & quercetin	0-5
15	Yang <i>et al.</i> (2017)	Hessian	37-45	Quercetin	0-0.6
16	Dogan (2018)	Lohmann Brown	40-48	Quercetin	0-20
17	Goliomytis <i>et al.</i> (2018)	Lohmann Brown	54-63	Hesperidin	0-1.5
18	Shahyar <i>et al.</i> (2018)	Hy-line Brown	58-70	Hesperitin	0-20
19	Simitzis <i>et al.</i> (2018)	Lohmann Brown	70-74	Quercetin	0-0.8
20	Abid & Ahmed (2019)	ISA Brown	40-50	Quercetin	0-1.2

Statistical analysis

The database was analyzed using a mixed-model methodology that has been widely used in meta-analysis research related to animal nutrition (Sauvant *et al.*, 2008; St-Pierre, 2001). Different studies were considered as random effects, and the doses of flavonoids administered were considered as fixed effects. The statistical model was as follows:

$$Y_{ij} = B_0 + B_1X_{ij} + B_2X_{ij}^2 + s_i + b_iX_{ij} + e_{ij}$$

where Y_{ij} = dependent variable, B_0 = overall intercept across all studies (fixed effect), B_1 = linear regression coefficient of Y on X (fixed effect), B_2 = quadratic regression coefficient of Y on X (fixed effect), X_{ij} = value of the continuous predictor variable (flavonoid addition levels), s_i = random effect of study i , b_i = random effect of study i on the regression coefficient of Y on X in study i , and e_{ij} = unexplained residual error. Linear and quadratic regression models were analyzed and considered significant effects for p -values below 5% ($P < 0.05$). The class statement was declared in the variable study because it does not contain any quantitative information. The p -value was used as the statistical model. All statistical analyses were carried out using R Software (version 3.60).

Results

The effects of flavonoids on the production performance of laying hens are presented in Table 2. Dietary addition of flavonoids linearly decreased ($P < 0.001$) feed intake of laying hens. The final body weight of laying hens was affected quadratically ($P < 0.001$) by flavonoids. The laying rate was not affected by the addition of flavonoids, while the FCR tended to decrease linearly ($P < 0.1$). Flavonoids positively affected egg quality profiles of laying hens (Table 3). The addition of flavonoids reduced egg and yolk cholesterol linearly (both at $P < 0.001$), and increased egg mass ($P < 0.001$, linearly), egg strength ($P < 0.05$, linearly) and yolk color ($P < 0.001$, linearly).

The effects of flavonoids on laying hens' liver and antioxidant enzyme profiles are presented in Tables 4 and 5, respectively. Triglyceride and cholesterol concentrations in the liver decreased linearly (at $P < 0.001$) after flavonoid supplementation, whereas liver weight was not affected. Superoxide dismutase was influenced quadratically ($P < 0.05$) following the addition of flavonoids, whereas catalase and glutathione peroxidase enzymes were not affected by dietary flavonoids supplementation.

Discussion

This meta-analysis generally indicates the potential of flavonoids as feed additives for substituting antibiotic growth promoters, which are now banned by most countries. Flavonoids do not impair egg production of laying hens and positively impact egg quality, liver

profile, and blood profile. Decreased feed intake due to flavonoids implies the greater feed efficiency in the laying hens. Such a decrease in feed intake has also been reported by Ariana *et al.* (2011), Deng *et al.* (2012) and Freitas *et al.* (2017). This decline is caused by the bitterness of flavonoids and thus reduced feed palatability (Goliomytis *et al.*, 2018). Flavonoids contribute to feed intake regulation even when supplemented at low levels (Wenk, 2002). Final body weight loss is associated with a decrease in feed intake and body fat reduction (Jennings *et al.*, 2017). The anti-lipogenic effect of flavonoids might be associated with their quercetin and hesperidin contents (Simitzis *et al.*, 2014; Zarrouki *et al.*, 2010). Quercetin plays a role in inhibiting cell population growth and adipogenesis in 3T3-L1 pre-adipocytes, which is a key to increasing fat mass. It seems that, at least in the short term, feed intake is not the primary factor affecting laying rate as reported by Oyedepi *et al.* (2007). In addition, flavonoids help the digestive tract to absorb nutrients by inhibiting pathogens and simultaneously increase beneficial genera such as *Bifidobacterium* and *Lactobacillus* (Pei *et al.*, 2020).

The decrease of triglycerides (in the liver) and cholesterol (both in liver and eggs) due to flavonoid supplementation was previously reported in laying hens (Kismiati *et al.*, 2020; Zhao *et al.*, 2013). Furthermore, flavonoid compounds such as quercetin, luteolin (Nekohashi *et al.*, 2014) and kaempferol (Salvamani *et al.*, 2014) help to decrease the cholesterol content of whole egg and yolk in laying hens. This may occur because flavonoids can form insoluble complexes with fat or cholesterol originating from feed, thereby inhibiting the absorption of endogenous and exogenous cholesterol (Zhao *et al.*, 2013; Kuang *et al.*, 2018; Lien *et al.*, 2008). Zhao *et al.* (2013) reported that triglycerides greatly influence the cholesterol content of eggs, which is in line with the results of the present study. Concerning reducing triglyceride levels, the antioxidant property of flavonoids stimulates lipoprotein-lipase activity (Roza *et al.*, 2007) that can hydrolyze VLDL, which converts triglyceride to fatty acids and glycerol. Flavonoids can increase superoxide dismutase due to the ability of flavonoids to resist oxidative disorders. Ion H^+ in flavonoid has donated an electron to anion superoxide to stabilize it and prevent DNA and lipoprotein oxidation (Wilmsen *et al.*, 2005; Aritanoga *et al.*, 2019).

The positive effect of flavonoids on egg mass is in line with Juráni *et al.* (2008) report, but in contrast with the study of Ahmad *et al.* (2017). Flavonoids provide a beneficial effect on the eggshell of laying hens. A decrease in FCR without affecting body weight indicates an increase in nutrient absorption, which was also reported by Prihambodo *et al.* (2020), mainly on calcium and phosphorus absorption.

Table 2. Regression equation on the influence of flavonoid (in g kg⁻¹ diet as fed) on growth performance of laying hens

Response Parameter	Unit	Model	N	Intercept	SE Intercept	Slope	SE Slope	P-value	RMSE	AIC	Trend
Final Body Weight	g	Q	22	1,670	36.4	-5.20	1.26	<0.001	0.981	215	Neg
Laying Rate	%	L	83	85.5	1.81	-0.118	0.094	0.212	1.44	512	Neg
Feed Conversion Rate	g	L	64	2.24	0.091	-0.007	0.004	0.078	1.47	1.75	Neg
Feed Intake	g/g	L	57	112	2.07	-0.369	0.090	<0.001	1.56	345	Neg

SE, standard error; RMSE, root mean square error; AIC, Akaike information criterion; Q, quadratic; L, linear; Min, minimal; Neg, negative.

^{b)}AIC is an estimator of the relative quality of statistical models for a given set of data (smaller is better)

Table 3. Regression equations on the influence of flavonoid (in g kg⁻¹ of diet as fed) on egg quality of laying hens

Response Parameter	Unit	Model	N	Intercept	SE Intercept	Slope	SE Slope	P-value	RMSE	AIC	Trend
Egg Weight	g	L	52	62.9	0.749	-0.002	0.026	0.941	1.65	198	Neg
Shape Index	%	L	17	75.5	0.413	0.019	0.090	0.945	1.06	52.6	Pos
Haugh Unit	Unit	L	76	82.9	1.24	0.048	0.108	0.657	1.40	392	Pos
Egg Mass	g	L	85	57.7	1.58	0.267	0.062	<0.001	1.52	397	Pos
Egg Thickness	10 ⁻² mm	L	73	39.3	1.22	0.091	0.066	0.173	1.91	340	Pos
Egg Strength	N	L	39	33.8	3.09	0.390	0.181	0.041	1.42	214	Pos
Egg Cholesterol	mg	L	29	184	14.6	-5.58	1.36	<0.001	0.97	257	Neg
Shell Weight	g	L	12	5.80	1.00	0.026	0.021	0.264	1.08	33.1	Pos
Shell Percentage	%	L	20	9.81	0.317	-0.001	0.016	0.935	1.19	44.2	Neg
Albumin Mass	g	L	16	18.4	8.96	0.004	0.031	0.894	1.31	70.9	Pos
Albumin Index	%	L	10	8.14	0.160	-0.063	0.246	0.808	0.98	17.8	Neg
Albumin Height	mm	L	10	6.65	0.751	0.002	0.096	0.982	1.08	35.0	Pos
Yolk Color	Unit	L	35	9.62	0.672	0.146	0.034	<0.001	1.21	84.2	Pos
Yolk Mass	g	L	78	15.9	0.468	0.027	0.019	0.169	1.52	176	Pos
Yolk: Egg Mass	g/g	L	65	26.1	0.661	-0.115	0.102	0.266	1.63	211	Neg
Yolk Cholesterol	mg/g	L	80	12.8	0.852	-0.296	0.052	<0.001	2.09	324	Neg

SE, standard error; RMSE, root mean square error; AIC, Akaike information criterion; L, linear; Pos, positive; Neg, negative

^{b)}AIC is an estimator of the relative quality of statistical models for a given set of data (smaller is better)

Table 4. Regression equations on the influence of flavonoid (in g kg⁻¹ of diet as fed) on liver profile of laying hens

Response Parameter	Unit	Model	N	Intercept	SE Intercept	Slope	SE Slope	P-value	RMSE	AIC	Trend
Liver Weight	g/100g	L	16	1.63	0.018	-0.007	0.014	0.606	1.06	-40.7	Neg
Liver Triglyceride	mg/g	L	16	14.8	0.243	-1.29	0.199	<0.001	1.06	32.7	Neg
Liver Cholesterol	mg/g	L	16	3.39	0.050	-0.205	0.038	<0.001	1.08	-12.8	Neg

SE, standard error; RMSE, root mean square error; AIC, Akaike information criterion; L, linear; Neg, negative

¹⁾AIC is an estimator of the relative quality of statistical models for a given set of data (smaller is better)

Table 5. Regression equations on the influence of flavonoid (in g kg⁻¹ of diet as fed) on antioxidant enzyme profile of laying hens

Response Parameter	Unit	Model	N	Intercept	SE Intercept	Slope	SE Slope	P-value	RMSE	AIC	Trend
Superoxide Dismutase	Unit	Q	16	28.2	2.23	1.72	1.39	0.024	1.03	115	Pos
Catalase	Unit	L	16	85.5	2.31	0.82	1.10	0.470	1.06	108	Pos
Glutathione Peroxidase	Unit	L	16	21.7	1.41	0.68	0.684	0.346	1.41	95.4	Pos

SE, standard error; RMSE, root mean square error; AIC, Akaike information criterion; Q, quadratic; L, linear; Max, maximum; Pos, positive

¹⁾AIC is an estimator of the relative quality of statistical models for a given set of data (smaller is better)

This also increases egg mass because shells are thicker and stronger. The presence of flavonoids also improved yolk color. As natural coloring agents in plants, flavonoids increase pigment deposition in egg yolks (Freitas *et al.*, 2017), making them look more reddish-yellow than ordinary eggs (Omri *et al.*, 2019).

Conclusion

Dietary supplementation of flavonoids is able to favourably modulate production performance and egg

quality of laying hens, and therefore reveals the potency of flavonoids for substituting antibiotic growth promoters for future production systems of laying hens.

Acknowledgements

This work is part of the research funded by the Indonesian Ministry of Research and Technology / National Research and Innovation Agency through the PMDSU grant.

References

- Abid AR & Ahmed SK. 2019. Influence of quercetin on some physiological measurements of layer hens. *Plant Archives*, 19(2): 3575-2582.
- Ahmad S, Khalique A, Pasha TN, Mehmood S, Hussain K, Ahmad S, Shaheen MS, Naeem M & Shafiq M. 2017. Effect of *Moringa oleifera* (Lam.) pods as feed additive on egg antioxidants, chemical composition and performance of commercial layers. *South African Journal of Animal Sciences*, 47: 864–874. DOI: 10.4314/sajas.v47i6.14
- Ariana M, Samie A, Edriss MA & Jahanian, R. 2011. Effects of powder and extract form of green tea and marigold, and α -tocopheryl acetate on performance, egg quality and egg yolk cholesterol levels of laying hens in late phase of production. *Journal of Medicinal Plants Research*, 5: 2710–2716. DOI: 10.5897/JMPR.9000537
- Aritanoga M, Effendi C & Herawati L. 2019. Gayo-arabica coffee decreases MDA and increases SOD after single bout submaximal physical exercise in Sedentary men. *Sumberdaya Hayati Journal*. 5: 58–63.
- Damaziak K, Riedel J, Gozdowski D, Niemiec J, Siennicka A & Rog D. 2017. Productive performance and egg quality of laying hens fed diets supplemented with garlic and onion extracts. *Journal of Applied Poultry Research*, 26: 337-349. DOI: 10.3382/japr/pfx001
- Deng W, Dong XF, Tong JM, Xie TH & Zhang Q. 2010. Effects of an aqueous alfalfa extract on production performance, egg quality and lipid metabolism of laying hens. *Journal of Animal Physiology and Animal Nutrition*, 96: 85–94. DOI: 10.1111/j.1439-0396.2010.01125.x
- Dogan C. 2018. The effects of licorice (*Glycyrrhiza glabra*) root on performance, some serum parameters and antioxidant capacity of laying hens. *Brazilian Journal of Poultry Science*, 20: 699-706. DOI: 10.1590/1806-9061-2018-0767
- El-Tarabany MS. 2018. Effect of royal jelly on behavioural patterns, feather quality, egg quality and some haematological parameters in laying hens at the late stage of production. *Journal of Animal Physiology Animal Nutrition*, 102: 599-606. DOI: 10.1111/jpn.12801
- Freitas ER, Fernandes DR, Souza DH, Dantas FDT, Santos RC, Oliveira GB, Cruz CEB, Braz NM, Câmara LF, Nascimento GAJ & Watanabe PH. 2017. Effect of syzygium cumini leaves on laying hens performance and egg quality. *Anais Da Academia Brasileira de Ciencias*, 89: 2479–2484. DOI: 10.1590/0001-3765201720150317
- Galal A, Abd El-Motaal AM, Ahmed AMH & Zaki TG. 2008. Productive performance and immune response of laying hens as affected by dietary propolis supplementation. *International Journal of Poultry Science*, 7: 272-278. DOI: 10.3923/jips.2008.272.278
- Goliomytis M, Orfanou H, Pertou E, Charismiadou MA, Simitzis PE & Deligeorgis SG. 2014. Effect of hesperidin dietary supplementation on hen performance, egg quality, and yolk oxidative stability. *British Poultry Science*, 55: 98-104. DOI: 10.1080/00071668.2013.870328
- Goliomytis M, Kostaki A, Avgoulas G, Lantzouraki, DZ, Siapi E, Zoumpoulakis P, Simitzis P & Deligeorgis SG. 2018. Dietary supplementation with orange pulp (*Citrus sinensis*) improves egg yolk oxidative stability in laying hens. *Animal Feed Science and Technology*, 244: 28–35. DOI: 10.1016/j.anifeedsci.2018.07.015
- Iskender H, Yenice G, Dokumacioglu E, Kaynar O, Hayirli A & Kaya A. 2017. Comparison of the effects of dietary supplementation of flavonoids on laying hen performance, egg quality and egg nutrient profile. *British Poultry Science*, 58: 550-556. DOI: 10.1080/00071668.2017.1349297
- Jennings A, MacGregor A, Spector T & Cassidy A. 2017. Higher dietary flavonoid intakes are

- associated with lower objectively measured body composition in women: evidence from discordant monozygotic twins. *The American Journal of Clinical Nutrition*, 105: 626-634. DOI: 10.3945/ajcn.116.144394
- Juráni M, Lamošsová D, MáčAjová M, Košťál L, Joubert E & Greksák M. 2008. Effect of rooibos tea (*Aspalathus linearis*) on Japanese quail growth, egg production and plasma metabolites. *British Poultry Science*, 49: 55–64. DOI: 10.1080/00071660701816949
- Kamboh AA, Arain MA, Mughal MJ, Zaman A & Arainl ZM. 2015. Flavonoids: Health promoting phytochemicals for animal production-a Review. *Journal of Animal Health and Production*, 3: 6–13. DOI: 10.1177/1461444810365020
- Kismiati S, Wahyuni HI, Muryani R, Sunarti D & Sumarsih S. 2020. Addition of binahong (*Anredera cordifolia*) leaf powder to diets to produce eggs with low cholesterol. *Veterinary World*, 13: 604–608. DOI: 10.14202/vetworld.2020.604-608
- Kuang H, Yang F, Zhang Y, Wang T & Chen G. 2018. The Impact of Egg Nutrient Composition and Its Consumption on Cholesterol Homeostasis. *Cholesterol*, 2. DOI: 10.1155/2018/6303810
- Kunatsa Y, Chidewe C & Zvidzai CJ. 2020. Phytochemical and anti-nutrient composite from selected marginalized Zimbabwean edible insects and vegetables. *Journal of Agriculture and Food Research*, 2. pp 100027. DOI: 10.1016/j.jafr.2020.100027
- Li XL, He WL, Yang ML, Yan YM, Xue YH, Xue YH & Zhao ST. 2017. Effect of dietary supplementation of *Ligustrum lucidum* on performance, egg quality and blood biochemical parameters of Hy-Line brown hens during the late laying period. *Animal*, 11: 1-6. DOI: 10.1017/S1751731117000532
- Lien TF, Yeh HS & Su WT. 2008. Effect of adding extracted hesperetin, naringenin and pectin on egg cholesterol, serum traits and antioxidant activity in laying hens. *Archives of Animal Nutrition*, 62: 33–43. DOI: 10.1080/17450390701780318
- Liu Y, Li Y, Liu HN, Suo YL, Hu LL, Feng XA, Zhang L & Jin F. 2013. Effect of quercetin on performance and egg quality during the laying period of hens. *British Poultry Science*, 54: 510-514. DOI: 10.1090/00071668.2013.799758
- Liu HN, Liu Y, Hu LL, Suo YL, Zhang L, Jin F, Feng XA, Teng N & Li Y. 2014. Effects of dietary supplementation of quercetin on performance, egg quality, cecal microflora populations, and antioxidant status in laying hens. *Poultry Science*, 93: 347-353. DOI: 10.3382/ps.2013-03225
- Lokhande A, Ingale SL, Lee SH, Sen S, Khong, Chae BJ & Kwon IK. 2014. Effects of dietary supplementation with *Gynura procumbens* (Merr.) on egg yolk cholesterol, excreta microflora and laying hen performance. *British Poultry Science*, 55: 524-531. DOI: 10.1080/00071668.2014.938020
- Nekohashi M, Ogawa M, Ogihara T, Nakazawa K, Kato H, Misaka T, Abe K & Kobayashi S. 2014. Luteolin and quercetin affect the cholesterol absorption mediated by epithelial cholesterol transporter Niemann-Pick C1-Like 1 in Caco-2 cells and rats. *PLoS ONE*, 9: 1–9. DOI: 10.1371/journal.pone.0097901
- Omri B, Alloui N, Durazzo A, Lucarini M, Aiello A, Romano R, Santini A & Abdouli H. 2019. Egg yolk antioxidants profiles: Effect of diet supplementation with linseeds and tomato-red pepper mixture before and after storage. *Foods*, 8. DOI: 10.3390/foods8080320
- Oyedeji JO, Orheruata AM & Omatsuli M. 2007. Effects of feed rationing on the laying performance of 40-weeks in-lay hens. *Journal of Food, Agriculture and Environment*, 5: 301–303.
- Panda V & Shinde P. 2017. Appetite suppressing effect of *Spinacia oleracea* in rats: Involvement of the short term satiety signal cholecystokinin. *Appetite*, 1: 224-230. DOI: 10.1016/k.appet.2017.02.030
- Pei R, Lui X & Bolling B. 2020. Flavonoids and gut health. *Current Opinion in Biotechnology*, 61: 153-159. DOI: 10.1016/j.copbio.2019.12.018
- Prihambodo TR, Nahrowi & Jayanegara A. 2019. Antibacterial Activity and Phytochemical Content of Silage Juice from Tropical Herbal Leaves. *IOP Conference Series: Materials Science and Engineering*, 546. DOI: 10.1088/1757-899X/546/4/042032
- Prihambodo TR, Sholikin MM, Qomariyah N, Jayanegara A, Baturbara I, Utomo DB & Ramli N. 2020. Effect of dietary flavonoids on performance, blood constituents, carcass composition and small intestinal morphology of broilers: A meta-analysis. *Asian-Australasian Journal of Animal Sciences*, 34: 434-442. DOI: 10.5713/ajas.20.0379
- Raina H, Soni G, Jauhari N, Sharma N & Bharadvaja N. 2014. Phytochemical importance of medicinal plants as potential sources of

- anticancer agents. *Turkish Journal of Botany*, 38: 1027-1035. DOI: 10.3906/bot-1405-93
- Roza JM, Xian-Liu Z & Guthrie N. 2007. Effect of citrus flavonoids and tocotrienols on serum cholesterol levels in hypercholesterolemic subjects. *Alternative Therapies in Health and Medicine*, 13: 44-48.
- Salvamani S, Gunasekaran B, Shaharuddin NA, Ahmad SA & Shukor MY. 2014. Antiatherosclerotic effects of plant flavonoids. *BioMed Research International*. DOI: 10.1155/2014/480258
- Sauvant D, Schmidely P, Daudin JJ & St-Pierre NR. 2008. Meta-analyses of experimental data in animal nutrition. *Animal*, 2: 1203–1214. DOI: 10.1017/S1751731108002280
- Shahyar HA, Ahmadzadeh A & Nobakht A. 2018. Effects of different levels of licorice (*Glycyrrhiza glabra*) medicinal plant powder on performance, egg quality and some of serum biochemical parameters in laying hens. *Iranian Journal of Applied Animal Science*, 8: 119-124.
- Simitzis P, Goliomytis M, Papalexis P, Veneti N, Charismiadou M, Kominakis A & Deligeorgis S. 2014. Effects of flavonoid dietary supplementation on egg yolk antioxidant capacity and cholesterol level. 65th Annual Meeting of the European Federation of Animal Science. 25-29 August. Copenhagen, Denmark.
- Simitzis P, Spanou D, Glasta N & Goliomytis M. 2018. Impact of dietary quercetin on laying hen performance, egg quality and yolk oxidative stability. *Animal Feed Science and Technology*, 239: 27-32. DOI: 10.1016/j.anifeedsci.2018.03.004
- St-Pierre NR. 2001. Invited review. Integrating quantitative findings from multiple studies using mixed model methodology. *Journal of Dairy Science*, 84: 741–755. DOI: 10.3168/jds.S0022-0302(01)74530-4
- Takahama U & Hirota S. 2018. Interactions of flavonoids with α -amylase and starch slowing down its digestion. *Food & Function*, 2: 677-687. DOI: 10.1039/c7fo01539a
- Ting S, Yeh HS & Lien TF. 2011. Effects of supplemental levels of hesperetin and naringenin on egg quality, serum traits and antioxidant activity of laying hens. *Animal Feed Science and Technology*, 163: 59-66. DOI: 10.1016/j.anifeedsci.2010.10.001
- Tugiyanti E & Susanti, E. 2018. The effect of breadfruit leaf flour (*Artocarpus altilis*) on number of blood cells and correlation between cholesterol blood and meat of Tegal ducks 10 weeks age. *Animal Production*, 19: 179. DOI: 10.20884/1.jap.2017.19.3.635
- Wenk C. 2003. Herbs and botanicals for monogastric animals. *Asian-Australian Journal of Animal Sciences*, 16: 282–290. DOI: 10.5713/ajas.2003.282
- Wilmsen PK, Spada DS & Salvador M. 2005. Antioxidant activity of the flavonoid hesperidin in chemical and biological systems. *Journal of Agricultural and Food Chemistry*, 53: 4757–4761. DOI: 10.1021/jf0502000
- Yang JX, Chaudhry MT, Yao JY, Wang SN, Zhou B, Wang M, Han CY, You Y & Li Y. 2017. Effects of phyto-oestrogen quercetin on productive performance, hormones, reproductive organs and apoptotic genes in laying hens. *Journal of Animal Physiology and Animal Nutrition*, 102: 505-513. DOI: 10.1111/jpn.12778
- Ying Y, Chun-yan H, Tabassum CM, Ling L, Jia-ying Y, Sheng-nan W, Jia-xin Y, Nan T & Yao L. 2015. Effect of quercetin on egg quality and components in laying hens of different weeks. *Journal of Northeast Agricultural University*, 22. L 23-32.
- Zarrouki B, Pillon NJ, Kalbacher E, Soula HA, Nia N'Jomen G, Grand L, Chambert S, Geloën A & Soulage CO. 2010. Cirsimarin, a potent antilipogenic flavonoid, decreases fat deposition in mice intra-abdominal adipose tissue. *International Journal of Obesity*, 34: 1566–1575. DOI: 10.1038/ijo.2010.85
- Zhao L, Zhang X, Cao F, Sun D, Wang T & Wang G. 2013. Effect of dietary supplementation with fermented Ginkgo-leaves on performance, egg quality, lipid metabolism and egg-yolk fatty acids composition in laying hens. *Livestock Science*, 155: 77–85. DOI: 10.1016/j.livsci.2013.03.024