



Validation of Prediction Equations of the Egg Characteristics in Laying Hens

Jhonatan Steve Inca¹  Diego Andres Martínez²  & Niceas Carlos Vilchez-Perales¹ 

¹ Department of Nutrition, Universidad Nacional Agraria La Molina, Lima, Peru

² Center of Excellence for Poultry Science, University of Arkansas, Fayetteville, AR 72701, USA

Poultry Science Journal 2022, 10(1): 71-82

Keywords

Laying
Equations
Validation
Egg characteristics

Corresponding author

Jhonatan Steve Inca
jhonastim7@gmail.com

Article history

Received: November 11, 2021
Revised: February 24, 2022
Accepted: February 25, 2022

Abstract

Poultry researchers have used mathematical models to explain some laying bird responses. There are many equations of the egg characteristics, but their validity has not yet been confirmed. Therefore, the aim of the current study was to assess the accuracy of the equations proposed to predict egg characteristics in laying hens. A total of one hundred forty-eight prediction equations of egg characteristics were collected from different studies. A total of 781 eggs from 75-week-old Hy-Line laying hens were gathered to measure egg quality characteristics, and other egg characteristics were calculated using these measurements. The residuals of the difference of observed and predicted values were used to calculate accuracy measurements like mean absolute deviation (MAD), mean squared error of prediction (MSEP), mean absolute percentage error (MAPE) and root mean squared error of prediction (RMSEP). RMSEP was used to estimate the error of the model (EM) with a 12% as the maximum level of validation of the egg characteristics prediction equation and 1.2% only for specific gravity. Nine egg characteristic prediction equations were validated with great accuracy because validated equations showed a value of MAD, MSEP, MAPE and RMSEP very low and EM less than 12%. Equations validated for external egg characteristic used easy-to-measure traits (i.e., egg weight, egg length, and egg width) as predictor variables. Fifteen egg characteristic prediction equations were validated with considerable accuracy. These equations might shorten the process of egg quality determination, reduce the waste of eggs, and thereby saving time and money.

Introduction

Egg production is regarded as one of the most profitable farm activities around the world because of the improvement of genetic and nutrition in laying hen. The majority of produced eggs are provided by conventional egg production systems (Matthews and Summer, 2015; USDA, 2018), and increasing the expenses related to the production process is of great concern to egg producers. Consumer acceptability and preferences are affected by characteristics so-called egg quality (Hisasaga *et al.*, 2020). In order to evaluate egg quality, thousands of eggs are wasted, lots of money is expensed (to buy materials and equipment), and so much time is lost (USDA, 2020). In addition, the assessment of egg quality is a complicated task since it requires, trained personal, specific materials and equipment, and an exclusive

place to efficiently conduct the complete process (Roberts *et al.*, 2013). Because egg quality determination is a complex process, making mistakes is more common. Hence, innovative alternatives are necessary to avoid measurement mistakes and to diminish egg production expenses (Inca, 2016).

Researchers are using mathematical models in a logical way to describe complex concepts, natural phenomenon, hypotheses and ideas. Models solve current issues using numerical solutions (Tedeschi and Menendez, 2020). Mathematical models are representations of the reality, and allow us to anticipate future behaviors outcomes (Tedeschi, 2006) or biological phenomena in animal science. In poultry science, many models have been developed to predict egg quality characteristics; but so far, the accuracy of those models has not been assessed.

Despite the fact that prediction models cannot entirely describe the reality under diverse conditions, they may contribute to observe some outcomes in a real system (Tedeschi, 2006). In a real system, a model will only have relevance if it represents the system with good accuracy (Hillston, 2003). Validation is a great form to verify if mathematical models are useful and accurate. Validation of models allows examining whether the intended model structure is appropriate -or the equation approximates the real system.

Many studies on laying birds have proposed mathematical models to estimate different egg quality characteristics. However, it has not been completely clarified if these models can predict egg quality characteristics with enough accuracy or they are useful enough to reduce the long procedure of egg quality characteristic determination. Therefore, the objective of the current study was to validate mathematical models to reduce mistakes and expenses in the egg quality determination process.

Materials and Methods

Biological materials and data collection

The experimental procedure was conducted at the Poultry Experimental Unit at the Universidad Nacional Agraria La Molina. Laying hens were raised in similar conditions according to Hy-line brown management recommendations. A total of 781 eggs from 75-week-age Hy-Line brown laying hens were collected during six weeks. At the end of the third week (77 weeks), egg quality characteristics were measured on every egg using a digital caliper (78-440, Stanley). Egg weight (EW, g), egg length (EL, mm), and egg width (EA, mm) were recorded. After cracking the eggs, shell weight (SW, g), shell thickness (ST, mm), albumen length (AL, mm; longest length), albumen width (AG, mm; perpendicular to AL), albumen height (AH, mm), yolk width (YG, mm; any direction), and yolk length (YL, mm; perpendicular to YG) were measured. The yolk was manually separated from the albumen and its weight (YW, g) and height (YH, mm) were determined. The albumen weight (AW, g) was determined with the following formula: $AW = EW - SW - YW$. Before cracking the eggs, the specific gravity (SG, g/cm^3) was determined using the saline solution technique as detailed by Inca *et al.* (2020). Furthermore, other external and internal characteristics (egg surface area, ESA; unit surface shell weight, U; shell ratio, SR; egg shape index, ESI; Haugh unit, HU; albumen index, AI; albumen ratio, AR; and yolk diameter, YD) were calculated by the calculations provided by Inca *et al.* (2020), as follows:

External egg characteristics:

Egg surface area (ESA, cm^2) = $3.9782 \times EW^{0.75056}$

Where, EW = egg weight (g).

Unit surface shell weight (U, mg/cm^2) = $\frac{SW}{ESA}$

Where, SW = shell weight (mg), ESA = egg surface area (cm^2).

Shell ratio (SR, %) = $\frac{SW}{EW} \times 100$

Where, SW= shell weight (mg), EW = egg weight(g).

Egg shape index (ESI, %) = $\frac{EG}{EL} \times 100$

Where, EG and EL are the egg width and length, respectively.

Internal egg characteristics:

Haugh unit (HU) = $100 \times \log (AH - 1.7 EW^{0.37} + 7.6)$

Where, AH = albumen height (mm), EW = egg weight (g).

Albumen index (AI, %) = $\frac{AH}{[AL+AG]/2} \times 100$

Where, AH = albumen height (mm), AL = albumen length (mm), AG = albumen width (mm).

Albumen ratio (AR, %) = $\frac{AW}{EW} \times 100$

Where, AW=albumen weight (g), EW=egg weight(g).

Yolk diameter (YD, mm) = $\frac{YG+YL}{2}$

Where, YG = yolk width (mm), YL = yolk length (mm).

Equations for validation

A total of 148 equations were compiled from various studies using the Cochrane method (Chandler and Hopewell, 2013). All equations pretended to predict different egg characteristics using simple or polynomial linear regression. Each corresponding determination coefficient (R^2) was collected. All prediction equations collected are shown in Table 1.

Validation procedure

First, predicted values were estimated using prediction equations of the egg characteristics. These values were compared with observed ones (measurements obtained in the current study) to obtain residuals of every egg characteristic, according to Tedeschi (2006). Second, residuals were used to calculate the error of every equation. In order to measure the prediction accuracy of the models, mean absolute deviation (MAD), mean squared error of prediction (MSEP), mean absolute percentage error (MAPE) and root mean squared error of prediction (RMSEP) were used. Formulas to determine the prediction accuracy of the models were based on Inca (2016) and Vaneput (2020) studies, as follows:

Mean absolute deviation (MAD) = $\frac{\sum_{i=1}^n |Y_i - \hat{Y}_i|}{n}$

Mean square error of prediction (MSEP) = $\frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n} = \frac{\sum_{i=1}^n E_i^2}{n}$

$$\text{Mean absolute percentage error (MAPE)} = \frac{\sum_{i=1}^n \left| \frac{Y_i - \hat{Y}_i}{Y_i} \right|}{n} = \frac{\sum_{i=1}^n \left| \frac{E_i}{Y_i} \right|}{n}$$

$$\text{Root mean squared error of prediction (RMSEP)} = \sqrt{\frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n}} = \sqrt{\frac{\sum_{i=1}^n E_i^2}{n}}$$

Where, Y_i = observed values of egg quality characteristic, \hat{Y}_i = predicted values of egg quality characteristic, E_i = sum of residuals, E_i^2 = sum of squared residuals, n = number of observations.

MAD outcomes were analyzed by one-way ANOVA, using GLM procedure. A comparison was made between MAD outcomes for each egg characteristic.

Differences between mean MAD values were determined using Tukey test at 0.05 significance level. All data were analyzed with RStudio software version 1.4.1106 (RStudio Team, 2016). Finally, the error of the model was estimated using RMSEPP (Salvador and Guevara, 2013) and with the following formula:

$$\text{Error of model (EM, \%)} = \frac{RMSEP}{\bar{Y}} \times 100$$

Where, RMSEP = root mean squared error of prediction, \bar{Y} = average value of the egg characteristics. For validation process, the maximum value established by the model of error was 12% for egg characteristics, and 1.2% only for specific gravity.

Table 1. Egg characteristics equations collected to be validated

N°	Reference	Year	Species	N*
R01	Nordstrom and Ousterhout	1982	Hen	14
R02	Harms <i>et al.</i>	1990	Hen	2
R03	Iposu <i>et al.</i>	1993	Hen	4
R04	Khurshid <i>et al.</i>	2003	Quail	16
R05	Seker	2004	Quail	6
R06	Narushin	2005	Hen	2
R07	Abanikannda and Leihgh	2007	Hen	6
R08	Yakubu <i>et al.</i>	2008	Chicken	17
R09	Alkan <i>et al.</i>	2008	Quail	10
R10	Fajemilehin	2008	Guinea Hen	18
R11	Alkan <i>et al.</i>	2009	Quail	28
R12	Callejo Ramos <i>et al.</i>	2010	Hen	5
R13	Çiçek <i>et al.</i>	2013	Quail	10
R14	Onunkwo and Okoro	2015	Guinea Hen	10

*N: Number of equations that each investigation provided.

Results

MAD, MSEP, MAPE and RMSEP and EM values from external egg characteristics are shown in Table 2. Egg weight equations were not accurate enough to be validated since all models had an EM greater than 12% (maximum value to validate equations). MAD, MSEP, MAPE, RMSEP and EM values indicated that egg length and egg width prediction models are not validated. These results showed no adequate accuracy, even when every collected equation had egg weight as a predictor variable. Moreover, all equations for egg length and egg width overestimated the values for prediction accuracy and EM, thus, none of the equations was validated (Table 2).

Both E31 and E40 indicated significant differences ($P < 0.05$) between shell weight equations for MAD. For these equations, MAD, MSEP and MAPE showed the lowest values and EM had a value

of 10.52% and 10.82%, respectively, which are reasons to validate E31 and E40. Moreover, E31 and E40 have the egg length and egg width as predictor variables; therefore, E31 and E40 are useful models because egg weight and egg width are easy characteristics to measure (Table 3). Twelve equations were collected to predict shell thickness, but none of them was validated, because they had an EM greater than 12%. On the other hand, E60 and E63 were validated for specific gravity since these models pointed a lowest MAD and were significantly different ($P < 0.05$) from the other models of specific gravity. In addition, MAD, MSEP and MAPE had the lowest values. Thus, these results showed an incredible accuracy of E60 and E63 (Table 3). E60 and E63 had EM of 1.03 and 0.70%, respectively, lower than 1.2% (established only for this egg characteristic).

Table 2. MAD, MSEP, MAPE and EM values of equations of the external egg characteristic

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Egg Weight						
E01	$Y = -21.658 + 0.828 X_A + 0.373 X_L (R^2 = 0.8260)$	R09	28.54 ^c	833.72	43.37	44.12
E02	$Y = 41.595 - 0.599 X_L (R^2 = 0.2110)$	R10	27.41 ^c	793.68	41.32	43.04
E03	$Y = 39.318 - 0.205 X_A (R^2 = 0.1682)$	R10	27.03 ^c	771.15	40.75	42.43
E04	$Y = -3.3133600 + 1.835144 X_L + 2.655127 X_A (R^2 = 0.3186)$	R04	46.18 ^e	2163.38	70.40	71.07
E05	$Y = 1.970096 + 2.252730 X_L (R^2 = 0.2313)$	R04	41.97 ^d	1791.18	63.92	64.66
E06	$Y = -1.0109318 + 3.616882 X_A (R^2 = 0.1781)$	R04	50.57 ^f	2591.51	77.12	77.78
E07	$Y = 14.89 + 5.40 X_L (R^2 = 0.2970)$	R08	18.44 ^b	366.06	27.70	29.23
E08	$Y = 22.21 + 5.34 X_A (R^2 = 0.3010)$	R08	19.77 ^b	423.16	29.66	31.43
E09	$Y = 14.71 + 3.10 X_L + 3.17 X_A (R^2 = 0.3444)$	R08	18.37 ^b	364.42	27.57	29.17
E10	$Y = 15.30 X_L^{0.63} (R^2 = 0.2958)$	R08	18.42 ^b	366.08	27.65	29.23
E11	$Y = 23.60 X_A^{0.44} (R^2 = 0.2863)$	R08	20.19 ^b	440.85	30.28	32.08
E12	$Y = 2.3887 + 1.3179 X_p (R^2 = 0.8850)$	R05	10.88 ^a	125.07	16.79	17.09
E13	$Y = 7.5927 - (0.2424 X_p) + 0.1160 X_p^2 (R^2 = 0.8890)$	R05	117.56 ^h	15623.36	175.76	190.98
E14	$Y = 3.1954 + 2.1929 X_y (R^2 = 0.5930)$	R05	25.91 ^c	700.69	39.35	40.44
E15	$Y = 12.6076 - (2.9558 X_y) + 0.6995 X_y^2 (R^2 = 0.6060)$	R05	93.08 ^g	10168.05	141.44	154.07
E16	$Y = 5.9338 + 6.3427 X_s (R^2 = 0.3680)$	R05	20.49 ^b	453.00	31.01	32.52
E17	$Y = 11.5143 - (6.8720 X_s) + 7.7518 X_s^2 (R^2 = 0.3760)$	R05	201.36 ⁱ	44346.74	306.73	321.75
Egg length						
E18	$Y = 0.292 + 1.271 X_w (R^2 = 0.5320)$	R11	24.00 ^d	610.47	40.17	41.55
E19	$Y = 0.436 + 1.055 X_w (R^2 = 0.3620)$	R11	10.02 ^b	121.20	16.69	18.51
E20	$Y = 0.323 + 1.165 X_w (R^2 = 0.3880)$	R11	17.09 ^c	319.61	28.57	30.06
E21	$Y = 0.014 + 1.148 X_w (R^2 = 0.1990)$	R11	8.82 ^a	77.93	14.82	14.84
Egg Width						
E22	$Y = 0.239 + 1.056 X_w (R^2 = 0.7710)$	R11	25.41 ^a	675.57	57.56	59.15
E23	$Y = 0.244 + 1.136 X_w (R^2 = 0.5010)$	R11	30.65 ^c	975.16	69.47	71.07
E24	$Y = 0.204 + 1.188 X_w (R^2 = 0.2750)$	R11	34.02 ^d	1196.67	77.11	78.73
E25	$Y = 0.016 + 1.104 X_w (R^2 = 0.1550)$	R11	28.33 ^b	835.81	64.19	65.80

^{a-j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴ EM = Error of model. X_w = Egg weight, X_L = Egg length, X_A = Egg width, X_p = Albumen Weight, X_y = Yolk Weight, X_s = Shell Weight.

In the current study, results of equations of the external egg characteristics are presented in Table 4. There were significant differences ($P < 0.05$) in MAD between equations to estimate egg surface area so that E64 and E65 showed lower values of MSEP and MAPE. Also, EM was lower than 12% in E64 and E65 equations (10.11 and 6.29, respectively). Equations for unit surface shell weight (also called SWUSA; Zhang *et al.*, 2020) and shell ratio were not

validated because prediction errors had higher values and EM were greater than 12%. There were significant differences in MAD for egg shape index prediction equations. E76 showed a lower value for MSEP and MAPE, and had a 7.08 for EM value; therefore, E76 was validated. Equations validated for external egg characteristic used easy-to-measure traits (i.e., egg weight, egg length, and egg width) as predictor variables.

Table 3. MAD, MSEP, MAPE and EM values of equations of the egg shell characteristics

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
E26	$Y = 3.716 + 0.04561 X_L^5 + 0.170 X_A^6 - 0.00831 X_W^7$ ($R^2 = 0.0880$)	R10	0.96 ^f	1.25	14.99	18.21
E27	$Y = 3.789 + 0.07572 X_L$ ($R^2 = 0.0740$)	R10	1.92 ⁱ	4.19	30.20	33.28
E28	$Y = 3.587 + 0.179 X_A$ ($R^2 = 0.0820$)	R10	1.79 ^h	3.69	28.05	31.24
E29	$Y = 4.549 - 0.0101 X_W$ ($R^2 = 0.0840$)	R10	1.02 ^f	1.39	15.88	19.15
E30	$Y = 2.77 + 0.49 X_A$ ($R^2 = 0.0286$)	R08	1.27 ^g	2.02	19.77	23.09
E31	$Y = 2.15 + 0.06 X_W$ ($R^2 = 0.0390$)	R08	0.50 ^a	0.42	8.41	10.52
E32	$Y = 2.13 + 0.05 X_W + 0.007 X_L$ ($R^2 = 0.0290$)	R08	0.81 ^e	0.92	12.79	15.56
E33	$Y = 1.82 + 0.04 X_W + 0.26 X_A$ ($R^2 = 0.0372$)	R08	0.72 ^{cd}	0.75	11.36	14.05
E34	$Y = 2.79 X_A^{0.37}$ ($R^2 = 0.0338$)	R08	1.36 ^g	2.27	21.18	24.53
E35	$Y = 0.74 X_W^{0.48}$ ($R^2 = 0.0391$)	R08	0.78 ^{de}	0.85	12.19	14.97
E36	$Y = 0.573 + 0.01532 X_L + 0.0238 X_W$ ($R^2 = 0.5070$)	R09	3.11 ^j	10.12	49.96	51.73
E37	$Y = (2.0341 \times X_W) - 2.1014 \times (X_W / X_G^8)$ ($R^2 = 0.9230$)	R02	0.87 ^f	0.91	13.69	15.48
E38	$Y = -0.521102 + 0.310761 X_L + 0.4074 X_A$ ($R^2 = 0.0701$)	R04	3.03 ^j	9.69	48.65	50.61
E39	$Y = 0.138189 + 0.062933 X_W + 0.233078 X_L$ ($R^2 = 0.0785$)	R04	0.68 ^{bc}	0.69	10.90	13.50
E40	$Y = -0.001150 + 0.071568 X_W + 0.311496 X_A$ ($R^2 = 0.0710$)	R04	0.52 ^a	0.44	8.66	10.82
E41	$Y = 0.8626 + 0.06418 X_G$ ($R^2 = 0.4636$)	R01	5.35 ^j	29.13	84.94	87.76
E42	$Y = 4.8334 + 0.005217 X_W$ ($R^2 = 0.0045$)	R01	1.47 ^g	2.60	22.52	26.24
E43	$Y = 2.1205 + 0.0555 X_W$ ($R^2 = 0.4667$)	R01	0.61 ^b	0.57	9.84	12.29
E44	$Y = -5.9724 + 0.06586 X_G + 0.09906 X_W$ ($R^2 = 0.8902$)	R01	5.27 ^j	28.13	84.16	86.24
E45	$Y = 2.1205 + 0.05545 X_W$ ($R^2 = 0.4667$)	R01	0.62 ^b	0.57	9.87	12.32
Shell Thickness						
E46	$Y = 0.135 + 0.0031 X_L$ ($R^2 = 0.0540$)	R09	0.043 ^a	0.003	11.81	14.94
E47	$Y = 0.459 + 0.00367 X_L$ ($R^2 = 0.0310$)	R10	0.129 ^e	0.018	38.63	38.43
E48	$Y = 0.431 + 0.01451 X_A$ ($R^2 = 0.0410$)	R10	0.143 ^f	0.022	42.64	42.24
E49	$Y = 0.337 + 0.003608 X_W$ ($R^2 = 0.0350$)	R10	0.221 ^h	0.051	65.20	64.23
E50	$Y = 0.267 + 0.00376 X_L + 0.0146 X_A + 0.00375 X_W$ ($R^2 = 0.0380$)	R10	0.248 ⁱ	0.064	72.76	71.65
E51	$Y = 0.154646 + 0.076448 X_A$ ($R^2 = 0.0420$)	R04	0.139 ^{ef}	0.021	41.43	41.19
E52	$Y = 0.154721 + 0.000694 X_W + 0.073939 X_A$ ($R^2 = 0.0384$)	R04	0.173 ^g	0.032	51.36	50.72
E53	$Y = 11.9836 + 0.3057 X_G$ ($R^2 = 0.6073$)	R01	0.042 ^a	0.003	30.94	14.37
E54	$Y = 42.3934 - 0.0768 X_W$ ($R^2 = 0.0355$)	R01	0.036 ^a	0.002	11.22	13.11
E55	$Y = -11.0561 + 0.4349 X_G + 0.2112 X_W$ ($R^2 = 0.7667$)	R01	0.070 ^b	0.002	27.90	13.09
E56	$Y = -0.17 + 0.01 X_W$ ($R^2 = 0.4120$)	R08	0.134 ^{de}	0.023	39.92	43.15
E57	$Y = 0.001 X_W^{1.55}$ ($R^2 = 0.4416$)	R08	0.304 ^j	0.103	88.74	91.27
Specific Gravity						
E58	$Y = 1.047 + 0.0842 X_T$ ($R^2 = 0.1220$)	R12	0.012 ^b	0.0002	1.06	1.15
E59	$Y = 1.070 + 0.00049 X_C$ ($R^2 = 0.1300$)	R12	0.015 ^c	0.0003	1.37	1.49
E60	$Y = 1.060 + 0.0028 X_H$ ($R^2 = 0.2550$)	R12	0.009 ^a	0.0001	0.86	1.03
E61	$Y = 1.040 + 0.00044 X_U$ ($R^2 = 0.3020$)	R12	0.013 ^b	0.0002	1.21	1.37
E62	$Y = 1.023 + 0.0572 X_T + 0.000456 X_C + 0.00036 X_U$ ($R^2 = 0.4530$)	R12	0.013 ^b	0.0002	1.22	1.33
E63	$Y = (X_W/0.9680) \times (X_W - X_S) + (0.4921 \times X_S)$ ($R^2 = 0.9230$)	R02	0.007 ^a	0.0001	0.64	0.70

^{a-j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴ EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width, X_T = Shell thickness, X_G = Specific gravity, X_S = Shell weight, X_U = Haugh unit, X_C = Yolk color, X_H = Albumen height.

Table 4. MAD, MSEP, MAPE and EM values of equations of the external egg characteristics

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Egg Surface Area						
E64	$Y = 10.561 - 0.178 X_A - 0.045 X_L + 1.535 X_W$ ($R^2 = 0.9960$)	R09	8.85 ^b	85.88	9.49	10.12
E65	$Y = 6.254 + 1.387 X_W$ ($R^2 = 0.9940$)	R09	5.36 ^a	33.28	5.71	6.29
E66	$Y = (0.9658 X_A / X_L + 2.1378) X_L * X_A$ ($R^2 = 0.9610$)	R06	62.31 ^c	3936.06	68.89	68.44
E67	$Y = (3.155 - 0.0136 X_L + 0.0115 X_A) X_L \times X_A$ ($R^2 = 0.9610$)	R06	62.25 ^c	3936.51	68.87	68.44
Unit Surface Shell Weight						
E68	$Y = 19.1936 + 0.6767 X_G$ ($R^2 = 0.6561$)	R01	45.31 ^a	2092.99	69.13	68.11
E69	$Y = 8.6878 - 0.0176 X_W$ ($R^2 = 0.0412$)	R01	59.63 ^b	3606.66	88.65	89.41
E70	$Y = -30.6697 + 0.9565 X_G + 0.4571 X_W$ ($R^2 = 0.8207$)	R01	62.96 ^c	4008.40	96.49	94.26
Shell Ratio						
E71	$Y = 12.49 - 0.06 X_W$ ($R^2 = 0.2400$)	R03	1.10 ^b	1.75	11.31	14.01
E72	$Y = -0.3773 + 0.1140 X_G$ ($R^2 = 0.8751$)	R01	9.26 ^c	86.54	102.84	98.65
E73	$Y = 13.4811 - 0.0728 X_W$ ($R^2 = 0.3295$)	R01	1.01 ^a	1.52	10.49	13.05
E74	$Y = -0.8794 + 0.1169 X_G + 0.004604 X_W$ ($R^2 = 0.8758$)	R01	9.44 ^c	89.87	104.83	100.53
Egg Shape Index						
E75	$Y = 0.78 - 0.00048 X_W + 0.0311 X_A - 0.0241 X_L$ ($R^2 = 0.9880$)	R09	5.87 ^b	44.23	8.12	8.99
E76	$Y = 0.79 + 0.0307 X_A - 0.02423 X_L$ ($R^2 = 0.9880$)	R09	4.34 ^a	27.46	6.04	7.08

^{a-j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴ EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width, X_G = Specific gravity.

Prediction errors and EM values of prediction equations of the albumen characteristics are shown in Table 5. Results showed that E91 predicted albumen weight with great accuracy, because MAD was significantly different ($P < 0.05$) rather than other equations, and the value of MSEP and MAPE were lower and EM had a value of 7.15% (lower than 12%). These outcomes validated E91 equation. Equations collected to predict albumen length and albumen height had not enough accuracy to be validated. However, two models of albumen width were validated, E100 and E102 presented smaller values for prediction errors. Moreover, MAD values had a significant different ($P < 0.05$) in comparison with other models and EM values were less than 12%.

Table 6 presents prediction errors and errors of model values of prediction equations of the yolk characteristics. Only E124 was validated for yolk weight because MAD value showed a significant

difference ($P < 0.05$) compared to other models. Additionally, MSEP and MAPE values were lower and EM was under 12%. The E134 had great prediction accuracy of yolk height. Prediction errors presented lower values and EM value was below 12% for E134. Equations to predict yolk width were not validated.

None of the equations was validated to predict the Haugh unit, yolk index and yolk diameter (Table 7). E146 was validated to predict albumen ratio. E146 showed a MAD value significantly different than E145 ($P < 0.05$), MSEP and MAPE values were lower, and EM had a value of 8.8.

A total of 15 equations were validated with reasonable accuracy (Table 8). These equations describe shell weight ($n=2$), specific gravity ($n=2$), egg surface area ($n=2$), egg shape index ($n=2$), albumen weight ($n=1$), albumen width ($n=2$), yolk weight ($n=1$), yolk height ($n=1$) and albumen ratio ($n=2$).

Table 5. MAD, MSEP, MAPE and EM values of equations of the albumen characteristics

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Albumen Weight						
E77	$Y = -1.214 + 1.185 X_W (R^2 = 0.8210)$	R07	36.75 ^m	1361.38	93.92	93.20
E78	$Y = -1.185 + 1.172 X_W (R^2 = 0.6740)$	R07	35.93 ^l	1301.22	91.83	91.12
E79	$Y = -0.456 + 0.989 X_W (R^2 = 0.6110)$	R07	24.68 ⁱ	614.91	63.28	62.64
E80	$Y = -1.428 + 1.223 X_W (R^2 = 0.7440)$	R07	39.02 ⁿ	1534.94	99.69	98.96
E81	$Y = 0.314 + 0.803 X_W (R^2 = 0.3970)$	R07	13.28 ^g	180.16	34.33	33.90
E82	$Y = -0.533 + 1.009 X_W (R^2 = 0.6720)$	R07	25.91 ^j	677.61	66.41	65.75
E83	$Y = -0.686 + 0.461 X_W + 0.0798 X_L + 0.412 X_A (R^2 = 0.7271)$	R04	7.88 ^{cd}	68.88	19.47	20.96
E84	$Y = -0.553150 + 0.468198 X_W + 0.426649 X_A (R^2 = 0.7275)$	R04	7.67 ^c	65.41	18.93	20.43
E85	$Y = 0.279557 + 0.468198 X_W (R^2 = 0.7224)$	R04	8.70 ^{de}	82.59	21.54	22.96
E86	$Y = -2.128934 + 0.925133 X_L + 1.63522 X_A (R^2 = 0.2919)$	R04	29.03 ^k	866.31	72.99	74.35
E87	$Y = -1.258 + 0.499 X_W (R^2 = 0.5460)$	R14	8.22 ^e	73.82	20.39	21.70
E88	$Y = 9.690 + 0.261 X_W (R^2 = 0.2100)$	R14	12.56 ^f	171.63	31.04	33.09
E89	$Y = 10.194 + 0.215 X_W (R^2 = 0.4320)$	R14	15.33 ^h	250.98	38.05	40.02
E90	$Y = -9.43 + 0.68 X_W (R^2 = 0.4260)$	R08	4.60 ^b	24.54	11.50	12.51
E91	$Y = 0.06 X_W^{1.54} (R^2 = 0.4382)$	R08	2.37 ^a	8.02	6.14	7.15
Albumen Length						
E92	$Y = -53.843 + 2.712 X_A + 1.376 X_L (R^2 = 0.2530)$	R09	59.48 ^f	3623.90	69.63	68.67
E93	$Y = 0.402 + 1.255 X_W (R^2 = 0.1870)$	R11	8.00 ^a	119.05	8.65	12.45
E94	$Y = 0.329 + 1.233 X_W (R^2 = 0.1760)$	R11	8.67 ^b	136.15	9.37	13.31
E95	$Y = 0.219 + 1.045 X_W (R^2 = 0.3010)$	R11	19.08 ^c	450.66	21.16	24.21
E96	$Y = 0.019 + 1.228 X_W (R^2 = 0.1890)$	R11	9.02 ^b	144.87	9.74	13.73
E97	$Y = -2.427 + 1.126 X_L (R^2 = 0.6281)$	R13	23.13 ^d	619.93	25.65	28.40
E98	$Y = 8.770 + 0.815 X_A (R^2 = 0.6280)$	R13	43.09 ^e	1963.29	48.52	50.54
Albumen Width						
E99	$Y = 0.877 + 0.599 X_W (R^2 = 0.4040)$	R11	35.88 ^d	1353.69	46.76	48.44
E100	$Y = 0.354 + 1.156 X_W (R^2 = 0.3880)$	R11	6.21 ^a	69.95	8.15	11.01
E101	$Y = 0.354 + 1.211 X_W (R^2 = 0.3560)$	R11	7.26 ^b	84.99	9.82	12.14
E102	$Y = 0.013 + 1.160 X_W (R^2 = 0.1620)$	R11	6.21 ^a	70.07	8.14	11.02
E103	$Y = -101.895 + 4.092 X_L (R^2 = 0.6825)$	R13	65.51 ^c	4414.60	88.02	87.47
E104	$Y = 5.035 + 0.864 X_A (R^2 = 0.6672)$	R13	32.97 ^c	1168.55	42.64	45.00
Albumen Height						
E105	$Y = -0.493 + 1.201 X_W (R^2 = 0.2850)$	R11	70.82 ⁱ	5076.66	1015.01	977.38
E106	$Y = 0.149 + 0.443 X_W (R^2 = 0.2870)$	R11	21.85 ^d	487.71	316.00	302.94
E107	$Y = -0.333 + 0.581 X_W (R^2 = 0.2650)$	R11	30.40 ^e	940.34	438.06	420.65
E108	$Y = -0.028 + 0.678 X_W (R^2 = 0.2220)$	R11	37.06 ^f	1394.20	533.03	512.20
E109	$Y = -16.774 + 1.003 X_W (R^2 = 0.6538)$	R10	41.58 ^h	1772.37	597.94	577.50
E110	$Y = -14.241 + 0.997 X_W - 0.464 X_A - 0.170 X_L (R^2 = 0.6844)$	R10	40.67 ^g	1695.78	584.94	564.88
E111	$Y = 24.236 - 0.690 X_A (R^2 = 0.3110)$	R10	13.92 ^b	195.35	202.30	191.73
E112	$Y = 29.097 - 0.836 X_L (R^2 = 0.3180)$	R10	16.84 ^c	285.02	243.72	231.59
E113	$Y = -0.326 + 0.196 X_W (R^2 = 0.3958)$	R13	5.21 ^a	30.66	78.47	75.96
E114	$Y = -0.484 + 0.196 X_W (R^2 = 0.3958)$	R13	5.05 ^a	29.04	76.21	73.92

^{a-j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴ EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width.

Table 6. MAD, MSEP, MAPE and EM values of equations of the yolk characteristics

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Yolk Weight						
E115	$Y = 10.525 + 0.01093 X_W + 0.295 X_L + 0.124 X_A$ ($R^2 = 0.0200$)	R10	3.08 ^b	12.67	17.71	21.47
E116	$Y = 12.224 + 0.007073 X_W$ ($R^2 = 0.6000$)	R10	3.91 ^c	18.78	22.69	26.14
E117	$Y = 11.708 + 0.161 X_A$ ($R^2 = 0.0850$)	R10	4.18 ^c	21.02	24.31	27.65
E118	$Y = 11.494 + 0.314 X_L$ ($R^2 = 0.1770$)	R10	3.25 ^b	13.93	18.73	22.51
E119	$Y = -0.618041 + 0.339520 X_W + 0.156591 X_A$ ($R^2 = 0.6080$)	R04	5.76 ^d	36.69	35.63	36.53
E120	$Y = -0.303204 + 0.355813 X_W$ ($R^2 = 0.6054$)	R04	6.45 ^e	45.29	39.81	40.59
E121	$Y = 0.003214 + 1.141682 X_A$ ($R^2 = 0.0830$)	R04	11.56 ^h	137.06	69.38	70.61
E122	$Y = 0.050845 + 0.921437 X_L$ ($R^2 = 0.1848$)	R04	11.05 ^g	125.40	66.28	67.54
E123	$Y = -3.663 + 0.405 X_W$ ($R^2 = 0.5830$)	R14	6.31 ^e	44.38	38.86	40.18
E124	$Y = 3.358 + 0.214 X_W$ ($R^2 = 0.2050$)	R14	1.37 ^a	3.44	8.62	11.19
E125	$Y = 1.876 + 0.273 X_W$ ($R^2 = 0.6860$)	R14	3.27 ^b	13.24	20.46	21.95
E126	$Y = -1.37 + 0.43 X_W$ ($R^2 = 0.4773$)	R08	10.21 ^f	109.57	62.68	63.13
E127	$Y = 0.23 X_W^{1.14}$ ($R^2 = 0.4989$)	R08	10.49 ^f	116.28	64.28	65.04
Yolk Width						
E128	$Y = 0.543 + 0.811 X_W$ ($R^2 = 0.6530$)	R11	13.54 ^b	205.95	34.05	35.81
E129	$Y = 0.326 + 1.027 X_W$ ($R^2 = 0.2790$)	R11	27.46 ^c	789.98	68.85	70.13
E130	$Y = 0.202 + 1.181 X_W$ ($R^2 = 0.1250$)	R11	37.41 ^d	1447.58	93.73	94.93
E131	$Y = 0.015 + 1.266 X_W$ ($R^2 = 0.1820$)	R11	42.79 ^e	1886.09	107.17	108.36
E132	$Y = 11.706 - 0.136 X_L$ ($R^2 = 0.3541$)	R13	36.46 ^d	1336.40	90.91	91.21
E133	$Y = 2.152 + 0.458 X_W$ ($R^2 = 0.3537$)	R13	8.04 ^a	72.17	19.96	21.20
Yolk Height						
E134	$Y = 2.908 + 0.261 X_A$ ($R^2 = 0.4650$)	R09	0.96 ^a	1.34	6.18	7.62
E135	$Y = 0.354 + 0.675 X_W$ ($R^2 = 0.3580$)	R11	29.36 ^f	877.96	193.87	195.32
E136	$Y = 0.371 + 0.663 X_W$ ($R^2 = 0.1960$)	R11	28.59 ^e	832.84	188.80	190.24
E137	$Y = 0.236 + 0.788 X_W$ ($R^2 = 0.1810$)	R11	36.63 ^g	1364.35	241.89	243.49
E138	$Y = -0.021 + 1.094 X_W$ ($R^2 = 0.2440$)	R11	56.40 ^h	3225.47	372.35	374.38
E139	$Y = 7.848 + 0.140 X_W$ ($R^2 = 0.1470$)	R14	1.86 ^b	4.36	12.52	13.76
E140	$Y = 7.184 + 0.150 X_W$ ($R^2 = 0.1850$)	R14	1.86 ^b	4.40	12.48	13.82
E141	$Y = -2.018 + 0.372 X_W$ ($R^2 = 0.2924$)	R13	7.15 ^c	55.92	47.32	49.30
E142	$Y = 4.262 + 0.355 X_W$ ($R^2 = 0.2912$)	R13	12.32 ^d	156.14	81.51	82.37

^{a-j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴ EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width.

Table 7. MAD, MSEP, MAPE and EM values of equations of the internal egg calculated characteristics

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Haugh Unit						
E143	$Y = 117.74 - 0.85X_W (R^2 = 0.2350)$	R03	21.36 ^b	526.66	25.09	27.64
E144	$Y = 108.612 - 0.507X_W (R^2 = 0.1840)$	R14	10.30 ^a	139.55	12.38	14.23
Albumen Ratio						
E145	$Y = 58.06 + 0.12X_W (R^2 = 0.2500)$	R03	5.65 ^b	40.74	9.69	10.58
E146	$Y = 49.03 + 0.24X_W (R^2 = 0.3560)$	R03	4.53 ^a	28.40	7.79	8.83
Yolk Index						
E147	$Y = 0.164 + 0.006X_W (R^2 = 0.2120)$	R14	18.72	371.56	51.37	52.18
Yolk Diameter						
E148	$Y = -4.406 + 0.824X_A + 0.195X_L + 0.168X_W (R^2 = 0.7940)$	R09	13.25	181.47	32.38	32.74

^{a,j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error.

⁴ EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width.

Table 8. Equations validated of egg characteristics.

N	Equation	Reference	MAD ¹	MSEP ²	MAPE ³ (%)	EM ⁴ (%)
Shell Weight						
E31	$Y = 2.15 + 0.06 X_W (R^2 = 0.0390)$	R08	0.50 ^a	0.42	8.41	10.52
E40	$Y = -0.001150 + 0.071568 X_W + 0.311496 X_A (R^2 = 0.0710)$	R04	0.52 ^a	0.44	8.66	10.82
Specific Gravity						
E60	$Y = 1.060 + 0.0028 X_H (R^2 = 0.2550)$	R12	0.009 ^a	0.0001	0.86	1.03
E63	$Y = (X_W/0.9680) \times (X_W - X_S) + (0.4921 \times X_S) (R^2 = 0.0007)$	R02	0.007 ^a	0.0001	0.64	0.70
Egg Surface Area						
E64	$Y = 10.561 - 0.178 X_A - 0.045 X_L + 1.535 X_W (R^2 = 0.9960)$	R09	8.85 ^b	85.88	9.49	10.11
E65	$Y = 6.254 + 1.387 X_W (R^2 = 0.9940)$	R09	5.36 ^a	33.28	5.71	6.29
Egg Shape Index						
E75	$Y = 0.78 - 0.00048 X_W + 0.0311 X_A - 0.0241 X_L (R^2 = 0.9880)$	R09	5.87 ^b	44.23	8.12	8.99
E76	$Y = 0.79 + 0.0307 X_A - 0.02423 X_L (R^2 = 0.9880)$	R09	4.34 ^a	27.46	6.04	7.08
Albumen Weight						
E91	$Y = 0.06 X_W^{1.54} (R^2 = 0.4382)$	R08	2.37 ^a	8.02	6.14	7.15
Albumen Width						
E100	$Y = 0.354 + 1.156 X_W (R^2 = 0.3880)$	R11	6.21 ^a	69.95	8.15	11.01
E102	$Y = 0.013 + 1.160 X_W (R^2 = 0.1620)$	R11	6.21 ^a	70.07	8.14	11.02
Yolk Weight						
E124	$Y = 3.358 + 0.214 X_W (R^2 = 0.2050)$	R14	1.37 ^a	3.44	8.62	11.19
Yolk Height						
E134	$Y = 2.908 + 0.261 X_A (R^2 = 0.4650)$	R09	0.96 ^a	1.34	6.18	7.62
Albumen Ratio						
E145	$Y = 58.06 + 0.12X_W (R^2 = 0.2500)$	R03	5.65 ^b	40.74	9.69	10.58
E146	$Y = 49.03 + 0.24X_W (R^2 = 0.3560)$	R03	4.53 ^a	28.40	7.79	8.83

^{a,j} Means within the same column with different superscript letters are significantly different ($P < 0.05$).

¹ MAD = Mean absolute deviation. ² MSEP = Mean squared error of prediction. ³ MAPE = Mean absolute percentage error. ⁴

EM = Error of model. X_W = Egg weight, X_L = Egg length, X_A = Egg width, X_H = Albumen height, X_S = Shell weight.

Discussion

Results of egg weight prediction models are inconsistent, according to the determination coefficient; for instance, E13 had a high value of the determination coefficient ($R^2 = 0.8890$) that was greater than E12 ($R^2 = 0.8850$). E13 was supposed to predict egg weight with reasonable accuracy, but in the current study E13 showed the lowest prediction capacity of egg weight. Moreover, E21 was close to be validated even when its determination coefficient showed a lowest value ($R^2 = 0.1990$) compared to the other models. Moreover, egg weight did not have the ability to predict egg length and egg width with enough accuracy, even though Olawumi and Ogunlade (2008), Dermanović and Mitrović (2013), Ojedapo (2013), and Mube *et al.* (2014) indicate egg weight had a strong correlation with egg length and egg width.

Two models were validated for shell weight, this egg characteristic is highly related to shell strength, and egg producers use shell weight as an indicator of egg quality. These models may be helpful for egg industry because both models have egg weight and egg width as predictors and are easy to measure (Inca *et al.*, 2020). Shell thickness is an important indicator of egg break resistance and is highly correlated with the presence of cracks in shell. In the current investigation, it was not possible to validate any equation for shell thickness, but some models (i.e., E54 and E55) were close to being validated, E63 showed a result that agrees with its determination coefficient ($R^2 = 0.9230$) and is decent model to predict specific gravity, but E63 has shell weight as the predictor variable, and breaking egg is necessary to obtain this characteristic (Inca *et al.*, 2020). The same issue was presented with E58 and E61.

Although all the equations for egg surface area had higher determination coefficients (> 0.9610), just two equations were validated (E54 and E65). E65 has some advantages; for instance, egg weight is the only necessary and easy to measure variable. Also, E65 presented good accuracy to predict egg surface area. Likewise, two models were validated with enough accuracy for egg shape index, both concur with their determination coefficients ($R^2 = 0.9880$). Martínez (2005) mentioned that regression equations with two or more predictor variables estimate the response variable with appropriate accuracy.

Albumen weight is a good indicator of egg quality because it represents the most significant portion of the egg weight, compared to shell weight and yolk weight. Egg protein is mainly located in albumen (Sotherland and Rahn, 1987) containing about 10.5% protein (Willems *et al.*, 2013). In the current study, an equation was validated for albumen weight with great accuracy (E91). This equation has egg weight as a predictor variable, making E91 a valuable model for farm conditions. Also, the result is consistent with

Olawumi and Ogunlade (2008) and Alipanah *et al.* (2013) reports in which high phenotypic correlation was found between egg weight and albumen weight.

Albumen length equations were not validated, but E93 was almost near to predict with reasonable accuracy. Even though two models (E100 and E102) were validated for albumen width, the determination coefficients of both E100 and E102 are inconsistent with the results of the present study because these models showed very low R^2 compared with the other prediction equations of albumen width. According to R^2 of E100 and E102, the independent variable could not explain the dependent variable well (albumen width). This result is consistent with Martínez (2005) because he mentioned that the objective of regression analysis is not to get a high determination coefficient value, but to obtain accurate estimators of the regression coefficients values. Furthermore, prediction models for albumen height evidenced that predictor variables (egg weight, egg length and egg width) had no relation with response variable (albumen height). Alkan *et al.* (2008) and Alipanah *et al.* (2013) found low correlations between albumen height and egg weight, egg length and egg width, supporting the results in the current study.

The determination coefficient of E124 ($R^2 = 0.2050$) demonstrates a weak capacity to explain the total variation of the response variable (yolk weight). Previous studies have indicated that the exclusive use of the determination coefficient is not a reliable indicator to explain the total variation of the response variable (Achen, 1982; Goldberger, 1991; Hernández, 1993). In addition, E124 has egg weight as a predictor variable, making this equation useful to predict yolk weight. These results agree with Kul and Seker (2004) and Zhang *et al.* (2005), who reported the highest phenotypic correlation between egg weight and yolk weight. Equations for yolk width were far away to be validated. This result agrees with Fajemilehin *et al.* (2009), Onunkwo and Okoro (2015), and Inca *et al.* (2020), who reported low correlations between yolk width and egg weight. E134 was validated to predict yolk height with excellent accuracy; also, this model has egg width as a predictor variable, which is an easy measurement characteristic under farm conditions, making E134 a very appropriate model according to the aims of the current investigation.

E143 and E144 had not enough predictive accuracy in of Haught Unit, and both of them had egg weight as the predictor variable. Debnath and Ghosh (2015) indicated that egg weight has a low phenotypic correlation ($r = 0.19$) with Haught Unit in layers, meaning that egg weight is a poor predictor to estimate Haught Unit. E145 and E146 were validated to predict albumen ratio with good accuracy, even when both equations have a low determination coefficient (0.250 and 0.356, respectively). Once

again, results evidence that using the determination coefficient as the only index of regression models may generate mistakes (Martínez, 2005). Prediction equations for yolk index and yolk diameter were not accurate enough to be validated.

Most of the validated equations include, as predictors, egg characteristics that are easy to measure (egg weight, egg length and egg width). These prediction equations are very useful, because it is not necessary to break the egg in order to estimate its

characteristics. In addition, using the determination coefficient as an indicator of the relationship between the response and predictor variables is unreliable if it is used as the only indicator in regression equations. The equations validated in the present study may be used by egg producers, poultry researchers and animal nutritionists to investigate egg characteristics of different laying species because they can help to reduce the costs of egg quality characteristics determination.

References

- Abanikannda OT & Leigh AO. 2007. Allometric relationships between composition and size of chicken table eggs. *International Journal of Poultry Science* 6: 211-217. DOI: 10.3923/ijps.2007.211.217
- Achen CH. 1982. *Interpreting and using regression*. Sage publications, California. USA. Pages, 56-67.
- Alipanah M, Deljio J, Rokouie M & Mohammadnia, R. 2013. Heritability and genetic and phenotypic correlation of egg quality traits in Khazak layers. *Trakia Journal of Sciences*, 2: 175–180.
- Alkan S, Karabag K, Galic A & Balcioglu MS. 2008. Predicting yolk height, yolk width, albumen length, eggshell weight, egg shape index, eggshell thickness, egg surface area of Japanese quails using various eggs traits as regressors. *International Journal of Poultry Science*, 7: 85–88. DOI: 10.3923/ijps.2008.85.88
- Alkan S, Mendes M, Karabag K & Karsli T. 2009. Effects of selection on allometric relationships between egg components and egg weight in Japanese quails of different lines. *Archiv Fur Geflugelkunde*, 74: 121-125.
- Callejo Ramos A, Cardoso Maciel W, Daza Andrada A, Siqueira de Castro Teixeira R & Buxade Carbo C. 2010. Effect of bird age and storage system on physical properties of eggs from brown laying hens. *PUBVET, Publicações em Medicina veterinaria e Zootecnia*, 4: 961-71.
- Chandler J & Hopewell S. 2013. Cochrane Methods - twenty years experience in developing systematic review methods. *Systematic Reviews*, 2: 76. DOI: 10.1186/2046-4053-2-76
- Çiçek TR, Uckardes F, Narinc D & Aksoy T. 2013. Comparison of principal component regression with the least square method in prediction of internal egg quality characteristics in Japanese quails. *Kafkas Universitesi Veteriner Fakultesi Dergisi*, 17: 687-692.
- Debnath BC & Ghosh TK. 2015. Phenotypic correlations between some external and internal egg quality traits in Gramapriya layers. *Exploratory Animal and Medical Research*, 5: 78-85.
- Dermanović V & Mitrović S. 2013. Phenotype variability of external quality traits of fertile eggs of partridge Italian bread in semi extensive breeding system. *Biotechnology in Animal Husbandry*, 29: 477–482. DOI: 10.2298/BAH1303477D
- Fajemilehin SO. 2008. Predicting post-broken traits using the pre-broken traits as regressors in the eggs of helmeted guinea fowl. *African Journal of Agricultural Research*, 3: 578-580.
- Fajemilehin SOK, Odubola OO, Fagbuaro SS & Akinyemi MO. 2009. Phenotypic correlations between some external and internal egg quality traits in the Nigerian helmeted guinea fowl, *Numida meleagris galeata pallas*. *Applied Tropical Agriculture*, 14: 102-108.
- Goldberger AS. 1991. *A course in econometrics*. Cambridge. Massachusettes: Harvard University Press. 432 Pages.
- Harms RH, Rossi AF, Sloan DR, Miles RD & Christmas RB. 1990. A method for estimating shell weight and correcting specific gravity for egg weight in eggshell quality studies. *Poultry Science*, 69: 48–52. DOI: 10.3382/ps.0690048
- Hernández AJ. 1993. Uso y abuso del coeficiente de determinación. *ESIC market*, 79: 77–92.
- Hillston J. 2003. *Model validation and verification*. Edinburgh: University of Edinburgh.
- Hisasaga C, Shinn SE & Tarrant KJ. 2020. Survey of egg quality in commercially available table eggs. *Poultry Science*, 99: 7202-7206. DOI: 10.1016/j.psj.2020.09.049
- Iposu SO, Onwuka CF & Eruvbetine D. 1993. The relationship between selected egg quality traits and egg size. *Nigerian Journal of Animal Production* 21: 156-160. DOI: 10.51791/njapv21i1.1175
- Inca JS. 2016. Validación de ecuaciones de predicción de las características de la calidad de huevo en gallinas de última fase de producción. Master thesis. Universidad Nacional Agraria La Molina, Lima, Peru. 295 Pages.
- Inca JS, Martinez DA & Vilchez C. 2020. Phenotypic correlation between external and internal egg quality characteristics in 85-week-old laying hens. *International Journal of Poultry Science*, 19: 346-355. DOI: 10.3923/ijps.2020.346.355
- Khurshid A, Farooq M, Durrani F, Sarbiland K & Chand N. 2003. Predicting egg weight, Shell

- weight, Shell thickness and hatching chick weight of Japanese quails using various egg traits as regressors. *International Journal of Poultry Science*, 2: 164-167. DOI: 10.3923/ijps.2003.164.167
- Kul S & Seker I. 2004. Phenotypic correlation between some external and internal egg quality traits in the Japanese quail (*Coturnix coturnix Japonica*). *International Journal of Poultry Science*, 3: 400-405. DOI: 10.3923/ijps.2004.400.405
- Martínez E. 2005. Errores frecuentes en la interpretación del coeficiente de determinación lineal. *Anuario Jurídico y Económico Escurialense*, 38: 315-332.
- Matthews WA & Summer DA. 2015. Effects of housing system on the costs of commercial egg production. *Poultry Science*, 94: 552-557. DOI: 10.3382/ps/peu011
- Mube HK, Kana JR, Tadondjou CD, Yemdjie DD, Manjeli Y & Tegua A. 2014. Laying performances and egg quality of local barred hens under improved conditions in Cameroon. *Journal of Applied Biosciences*, 74: 6157-6163. DOI: 10.4314/jab.v74i1.8
- Narushin VG. 2005. Egg geometry calculation using the measurements of length and Breadth. *Poultry Science*, 84: 482-484. DOI: 10.1093/ps/84.3.482
- Nordstrom JO & Ousterhout LE. 1982. Estimation of shell weight and shell thickness from egg specific gravity and egg weight. *Poultry Science*, 61: 1991-1995. DOI: 10.3382/ps.0611991
- Ojedapo LO. 2013. Phenotypic correlation between the external and internal egg quality traits of pharaoh quail reared in derived savanna zone of Nigeria. *Journal of Biology, Agriculture and Healthcare*, 3: 80-84.
- Olawumi SO & Ogunlade JT. 2008. Phenotypic correlations between some external and internal egg quality traits in the exotic Isa Brown layer breeders. *Asian Journal of Poultry Science*, 2: 30-35. DOI: 10.3923/ajpsaj.2008.30.35
- Onunkwo DN & Okoro IC. 2015. Predicting egg quality and egg production traits using egg weight and body weight respectively in three varieties of helmeted guinea fowls in humid tropics. *International Journal of Livestock Research*, 5: 111-121. DOI: 10.5455/ijlr.20150329124016
- Roberts JR, Chousalkar K & Samiullah S. 2013. Egg quality and age of laying hens: implications for product safety. *Animal Production Science*, 53: 1291-1297. DOI: 10.1071/AN12345
- RStudio Team. 2016. RStudio: Integrated Development for R. Boston, USA.
- Salvador E & Guevara V. 2013. Development and validation of a model to predict the optimum requirement of essential amino acids and performance in commercial laying hens. Departamento de producción animal, facultad de medicina veterinaria y zootecnia. Universidad Nacional San Luis Gonzaga, Ica. Perú. *Revista de Investigaciones Veterinarias del Perú*, 24: 264-276. DOI: 10.15381/rivep.v24i3.2574
- Seker I. 2004. Prediction of albumen weight, yolk weight, and shell weight as egg weight in Japanese quail eggs. *Uludag University Journal of the Faculty of Veterinary Medicine*, 23: 87-92.
- Sotherland PR & Rahn H. 1987. On the composition of bird eggs. *Condor*, 89: 48-65. DOI: 10.2307/1368759
- Tedeschi LO. 2006. Assessment of the adequacy of mathematical models. *Agricultural Systems*, 89: 225-247. DOI: 10.1016/j.agry.2005.11.004
- Tedeschi LO & Menendez HM. 2020. Mathematical modeling in animal production. In: *Animal Agriculture: Sustainability, Challenges and Innovations*. Elsevier Publisher. USA. Pages, 431-453. DOI: 10.1016/B978-0-12-817052-6.00025-2
- USDA U.S. Department of Agriculture. 2018. World agricultural supply and demand estimates. WASDE 574. USDA, Washington, D.C.
- USDA. 2020. America's Farmers: Resilient Throughout the COVID Pandemic. Available: <https://www.usda.gov/media/>
- Vandeput, N. 2020. Data science for supply chain forecasting. Second Edition. De Gruyter. 282 pages. DOI: 10.1515/9783110671124-201
- Willems E, Wang Y, Willemsepn H, Lesuisse J, Franssens L, Guo X, Koppenol A, Buyse J, Decuyper E & Everaert N. 2013. Partial albumen removal early during embryonic development of layer-type chickens has negative consequences on laying performance in adult life. *Poultry Science*, 92: 1905-1915. DOI: 10.3382/ps.2012-03003
- Yakubu A, Ogah DM & Barde RE. 2008. Productivity and egg quality characteristics of free range naked neck and normal feathered Nigerian indigenous chickens. *International Journal of Poultry Science*, 7: 579-585. DOI: 10.3923/ijps.2008.579.585
- Zhang C, Ning H, Xu Y, Hou C & Yang N. 2005. Heritabilities and Genetic and Phenotypic Correlations of Egg Quality Traits in Brown-Egg Dwarf Layers. *Poultry Science*, 84: 1209-1213. DOI: 10.1093/ps/84.8.1209
- Zhang B, Weil J, Guerra AB, Maharjan P, Hilton K, Suesuttajit N, Martinez D & Coon CN. 2020. Egg shell quality and bone status as affected by environmental temperature, Ca and non-phytate O intake and in vitro limestone solubility in Single-Comb White Leghorn hens. *International Journal of Poultry Science*, 19: 219-231. DOI: 10.3923/ijps.2020.219.231